

表 3017 PTH値(pg/mL) 原疾患別(血液透析濾過患者 透析回数週3回) PTH測定法:intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	714	803	961	1,058	1,162	1,116	1,148	1,052	1,005	2,258	28	1,369	838	479	273	81	120	139	231	230	15,065	231	15,296	192.28	195.43
(%)	(4.7)	(5.3)	(6.4)	(7.0)	(7.7)	(7.4)	(7.6)	(7.0)	(6.7)	(15.0)	(0.2)	(9.1)	(5.6)	(3.2)	(1.8)	(0.5)	(0.8)	(0.9)	(1.5)	(1.5)	(100.0)				
慢性腎盂腎炎	18	33	28	29	38	33	25	23	26	70	3	31	13	13	15	2	4	2	2	9	417	6	423	192.91	226.37
(%)	(4.3)	(7.9)	(6.7)	(7.0)	(9.1)	(7.9)	(6.0)	(5.5)	(6.2)	(16.8)	(0.7)	(7.4)	(3.1)	(3.1)	(3.6)	(0.5)	(1.0)	(0.5)	(0.5)	(2.2)	(100.0)				
急速進行性糸球体腎炎	18	17	29	22	22	26	28	25	25	43		18	14	8	6		4	2	3	3	313	6	319	177.65	224.65
(%)	(5.8)	(5.4)	(9.3)	(7.0)	(7.0)	(8.3)	(8.9)	(8.0)	(8.0)	(13.7)		(5.8)	(4.5)	(2.6)	(1.9)		(1.3)	(0.6)	(1.0)	(1.0)	(100.0)				
妊娠腎/妊娠中毒症	22	24	20	28	31	23	22	27	25	44	1	23	12	10	11	2	3	3	6	7	344	15	359	184.11	173.74
(%)	(6.4)	(7.0)	(5.8)	(8.1)	(9.0)	(6.7)	(6.4)	(7.8)	(7.3)	(12.8)	(0.3)	(6.7)	(3.5)	(2.9)	(3.2)	(0.6)	(0.9)	(0.9)	(1.7)	(2.0)	(100.0)				
その他の分類不能の腎炎	9	11	15	17	12	16	15	14	18	41	1	18	22	5	7		4	2	3	5	235	4	239	208.33	194.70
(%)	(3.8)	(4.7)	(6.4)	(7.2)	(5.1)	(6.8)	(6.4)	(6.0)	(7.7)	(17.4)	(0.4)	(7.7)	(9.4)	(2.1)	(3.0)		(1.7)	(0.9)	(1.3)	(2.1)	(100.0)				
多発性囊胞腎	65	89	107	102	103	132	135	110	106	249	3	162	93	56	31	9	12	11	33	21	1,629	19	1,648	195.47	186.31
(%)	(4.0)	(5.5)	(6.6)	(6.3)	(6.3)	(8.1)	(8.3)	(6.8)	(6.5)	(15.3)	(0.2)	(9.9)	(5.7)	(3.4)	(1.9)	(0.6)	(0.7)	(0.7)	(2.0)	(1.3)	(100.0)				
腎硬化症	117	197	237	252	264	282	244	230	253	504	4	305	162	104	51	12	19	13	41	32	3,323	50	3,373	177.52	157.10
(%)	(3.5)	(5.9)	(7.1)	(7.6)	(7.9)	(8.5)	(7.3)	(6.9)	(7.6)	(15.2)	(0.1)	(9.2)	(4.9)	(3.1)	(1.5)	(0.4)	(0.6)	(0.4)	(1.2)	(1.0)	(100.0)				
悪性高血圧	15	17	24	22	25	28	24	25	20	58	1	38	16	9	6	1	2	4	4	2	341	3	344	180.81	153.43
(%)	(4.4)	(5.0)	(7.0)	(6.5)	(7.3)	(8.2)	(7.0)	(7.3)	(5.9)	(17.0)	(0.3)	(11.1)	(4.7)	(2.6)	(1.8)	(0.3)	(0.6)	(1.2)	(1.2)	(0.6)	(100.0)				
糖尿病性腎症	553	903	988	1,177	1,238	1,211	1,202	1,045	921	2,215	23	1,317	733	397	226	64	94	81	138	98	14,624	275	14,899	171.64	142.68
(%)	(3.8)	(6.2)	(6.8)	(8.0)	(8.5)	(8.3)	(8.2)	(7.1)	(6.3)	(15.1)	(0.2)	(9.0)	(5.0)	(2.7)	(1.5)	(0.4)	(0.6)	(0.6)	(0.9)	(0.7)	(100.0)				
SLE腎炎	18	30	23	25	29	28	23	22	24	45		29	20	10	7	2	4	5	9	6	359	8	367	204.43	284.18
(%)	(5.0)	(8.4)	(6.4)	(7.0)	(8.1)	(7.8)	(6.4)	(6.1)	(6.7)	(12.5)		(8.1)	(5.6)	(2.8)	(1.9)	(0.6)	(1.1)	(1.4)	(2.5)	(1.7)	(100.0)				
アミロイド腎	3	6	3	1	6	3	3	5	5	9	1	5	2	2	1			1	1		57	1	58	172.18	130.08
(%)	(5.3)	(10.5)	(5.3)	(1.8)	(10.5)	(5.3)	(5.3)	(8.8)	(8.8)	(15.8)	(1.8)	(8.8)	(3.5)	(3.5)	(1.8)			(1.8)	(1.8)		(100.0)				
痛風腎	8	10	9	9	12	11	8	8	5	22		17	8	2	4	2	3			5	143	4	147	204.31	224.14
(%)	(5.6)	(7.0)	(6.3)	(6.3)	(8.4)	(7.7)	(5.6)	(5.6)	(3.5)	(15.4)		(11.9)	(5.6)	(1.4)	(2.8)	(1.4)	(2.1)			(3.5)	(100.0)				
先天性代謝異常による腎不全		3	1	4	2	5	5	1		4		3	5	2	4			1	2	1	43		43	247.79	207.84
(%)		(7.0)	(2.3)	(9.3)	(4.7)	(11.6)	(11.6)	(2.3)		(9.3)		(7.0)	(11.6)	(4.7)	(9.3)			(2.3)	(4.7)	(2.3)	(100.0)				
腎・尿路結核	2	2	1			2	3	1	1	5		1	1	2					1		22		22	187.17	157.38
(%)	(9.1)	(9.1)	(4.5)			(9.1)	(13.6)	(4.5)	(4.5)	(22.7)		(4.5)	(4.5)	(9.1)					(4.5)		(100.0)				
腎・尿路結石	4	4	5	2	2	3	4	6	4	15	1	9	1	3	1	1			2	2	69	1	70	220.69	224.09
(%)	(5.8)	(5.8)	(7.2)	(2.9)	(2.9)	(4.3)	(5.8)	(8.7)	(5.8)	(21.7)	(1.4)	(13.0)	(1.4)	(4.3)	(1.4)	(1.4)			(2.9)	(2.9)	(100.0)				
腎・尿路腫瘍	4	6	8	9	12	13	10	7	4	9		10	3	2	2	1	1		2	1	104		104	166.06	152.40
(%)	(3.8)	(5.8)	(7.7)	(8.7)	(11.5)	(12.5)	(9.6)	(6.7)	(3.8)	(8.7)		(9.6)	(2.9)	(1.9)	(1.9)	(1.0)	(1.0)		(1.9)	(1.0)	(100.0)				
閉塞性尿路障害	3	4	3	8	4	8	7	4	5	11		12	2	3	5			1	3	3	86	1	87	246.19	307.87
(%)	(3.5)	(4.7)	(3.5)	(9.3)	(4.7)	(9.3)	(8.1)	(4.7)	(5.8)	(12.8)		(14.0)	(2.3)	(3.5)	(5.8)			(1.2)	(3.5)	(3.5)	(100.0)				
骨髄腫	1	1	4	2	2	2	1	2	1	3		2	1	2					1		25		25	169.60	142.23
(%)	(4.0)	(4.0)	(16.0)	(8.0)	(8.0)	(8.0)	(4.0)	(8.0)	(4.0)	(12.0)		(8.0)	(4.0)	(8.0)					(4.0)		(100.0)				
腎形成不全	2	10	6	5	5	15	8	4	3	22		13	6	3	4	2	1	1	4	3	117	1	118	230.12	238.01
(%)	(1.7)	(8.5)	(5.1)	(4.3)	(4.3)	(12.8)	(6.8)	(3.4)	(2.6)	(18.8)		(11.1)	(5.1)	(2.6)	(3.4)	(1.7)	(0.9)	(0.9)	(3.4)	(2.6)	(100.0)				
不明	146	232	250	324	384	292	299	278	238	611	8	371	214	118	79	11	23	33	59	51	4,021	93	4,114	186.25	187.15
(%)	(3.8)	(5.8)	(6.2)	(8.1)	(9.5)	(7.3)	(7.4)	(6.9)	(5.9)	(15.2)	(0.2)	(9.2)	(5.3)	(2.9)	(2.0)	(0.3)	(0.6)	(0.8)	(1.5)	(1.3)	(100.0)				
再導入	16	24	27	26	27	31	34	33	34	48	1	38	29	15	12	2	3	5	11	8	424	4	428	207.33	193.89
(%)	(3.8)	(5.7)	(6.4)	(6.1)	(6.4)	(7.3)	(8.0)	(7.8)	(8.0)	(11.3)	(0.2)	(9.0)	(6.8)	(3.5)	(2.8)	(0.5)	(0.7)	(1.2)	(2.6)	(1.9)	(100.0)				
その他	39	60	60	88	72	75	72	81	74	144		86	54	31	22	4	4	6	12	12	996	11	1,007	184.96	198.84
(%)	(3.9)	(6.0)	(6.0)	(8.8)	(7.2)	(7.5)	(7.2)	(8.1)	(7.4)	(14.5)		(8.6)	(5.4)	(3.1)	(2.2)	(0.4)	(0.4)	(0.6)	(1.2)	(1.2)	(100.0)				
合計	1,777	2,486	2,809	3,210	3,452	3,355	3,320	3,003	2,797	6,430	75	3,877	2,249	1,276	767	196	301	310	568	499	42,757	733	43,490	183.79	176.85
(%)	(4.2)	(5.8)	(6.6)	(7.5)	(8.1)	(7.8)	(7.8)	(7.0)	(6.5)	(15.0)	(0.2)	(9.1)	(5.3)	(3.0)	(1.8)	(0.5)	(0.7)	(0.7)	(1.3)	(1.2)	(100.0)				
記載なし	1	2	1					1		1											6		6	79.67	83.12
(%)	(16.7)	(33.3)	(16.7)					(16.7)		(16.7)											(100.0)				
総計	1,778	2,488	2,810	3,210	3,452	3,355	3,320	3,004	2,797	6,431	75	3,877	2,249	1,276	767	196	301	310	568	499	42,763	733	43,496	183.77	176.85
(%)	(4.2)	(5.8)	(6.6)	(7.5)	(8.1)	(7.8)	(7.8)	(7.0)	(6.5)	(15.0)	(0.2)	(9.1)	(5.3)	(3.0)	(1.8)	(0.5)	(0.7)	(0.7)	(1.3)	(1.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2015年調査