

表 3030 PTH値(pg/mL) 原疾患別（腹膜透析患者）PTH測定法intact-PTH

原疾患	20未満	20～	40～	60～	80～	100～	120～	140～	160～	180～	240～	241～	300～	360～	420～	480～	500～	540～	600～	800～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	29	60	104	99	78	102	83	96	87	221	3	156	97	52	44	12	24	24	45	54	1,470	105	1,575	244.04	252.25
(%)	(2.0)	(4.1)	(7.1)	(6.7)	(5.3)	(6.9)	(5.6)	(6.5)	(5.9)	(15.0)	(0.2)	(10.6)	(6.6)	(3.5)	(3.0)	(0.8)	(1.6)	(1.6)	(3.1)	(3.7)	(100.0)				
慢性腎盂腎炎	1	4	4	4	2	2	3	2	2	6		7	6	2	2		1	2	1		51	7	58	218.85	157.72
(%)	(2.0)	(7.8)	(7.8)	(7.8)	(3.9)	(3.9)	(5.9)	(3.9)	(3.9)	(11.8)		(13.7)	(11.8)	(3.9)	(3.9)		(2.0)	(3.9)	(2.0)		(100.0)				
急速進行性糸球体腎炎	3	1	3	1	1	3	5	2		1		3	7	1	1	1					33	2	35	190.52	134.02
(%)	(9.1)	(3.0)	(9.1)	(3.0)	(3.0)	(9.1)	(15.2)	(6.1)		(3.0)		(9.1)	(21.2)	(3.0)	(3.0)	(3.0)					(100.0)				
妊娠腎/妊娠中毒症			1	2			1	3	1	2		2	1		1	2			2		18	1	19	280.83	203.66
(%)			(5.6)	(11.1)			(5.6)	(16.7)	(5.6)	(11.1)		(11.1)	(5.6)		(5.6)	(11.1)			(11.1)		(100.0)				
その他の分類不能の腎炎		3	1	2	2	7		3	1	5		3	5	3	1			1	1	3	41	1	42	275.43	268.61
(%)		(7.3)	(2.4)	(4.9)	(4.9)	(17.1)		(7.3)	(2.4)	(12.2)		(7.3)	(12.2)	(7.3)	(2.4)			(2.4)	(2.4)	(7.3)	(100.0)				
多発性囊胞腎	1	11	8	8	12	5	7	6	3	22		16	5	7	2	2	2	5	3	5	130	8	138	250.89	280.93
(%)	(0.8)	(8.5)	(6.2)	(6.2)	(9.2)	(3.8)	(5.4)	(4.6)	(2.3)	(16.9)		(12.3)	(3.8)	(5.4)	(1.5)	(1.5)	(1.5)	(3.8)	(2.3)	(3.8)	(100.0)				
腎硬化症	14	36	35	33	32	41	28	29	36	77	1	72	37	28	17	8	4	6	19	15	568	39	607	228.81	204.07
(%)	(2.5)	(6.3)	(6.2)	(5.8)	(5.6)	(7.2)	(4.9)	(5.1)	(6.3)	(13.6)	(0.2)	(12.7)	(6.5)	(4.9)	(3.0)	(1.4)	(0.7)	(1.1)	(3.3)	(2.6)	(100.0)				
悪性高血圧	1			4	3	3	5	5	4	9		7	4	4	1	1	1	1		1	54	2	56	233.50	165.10
(%)		(1.9)		(7.4)	(5.6)	(5.6)	(9.3)	(9.3)	(7.4)	(16.7)		(13.0)	(7.4)	(7.4)	(1.9)	(1.9)	(1.9)	(1.9)		(1.9)	(100.0)				
糖尿病性腎症	46	73	74	92	94	83	92	77	65	200	3	164	102	55	39	11	12	18	23	17	1,340	114	1,454	208.61	179.31
(%)	(3.4)	(5.4)	(5.5)	(6.9)	(7.0)	(6.2)	(6.9)	(5.7)	(4.9)	(14.9)	(0.2)	(12.2)	(7.6)	(4.1)	(2.9)	(0.8)	(0.9)	(1.3)	(1.7)	(1.3)	(100.0)				
SLE腎炎			1	1	3		2	1		2		3	1	2				2	1		19	2	21	263.47	194.43
(%)			(5.3)	(5.3)	(15.8)		(10.5)	(5.3)		(10.5)		(15.8)	(5.3)	(10.5)				(10.5)	(5.3)		(100.0)				
アミロイド腎	1				1			2	1	1			1								7		7	162.39	98.91
(%)		(14.3)			(14.3)			(28.6)	(14.3)	(14.3)			(14.3)								(100.0)				
痛風腎	1			1	1	3		2	2	4		3							1	1	19	2	21	234.96	229.34
(%)		(5.3)		(5.3)	(5.3)	(15.8)		(10.5)	(10.5)	(21.1)		(15.8)							(5.3)	(5.3)	(100.0)				
先天性代謝異常による腎不全				1				1		2					1		2				7	2	9	295.60	195.58
(%)				(14.3)				(14.3)		(28.6)					(14.3)		(28.6)				(100.0)				
腎・尿路結核					1																1		1	99.00	
(%)					(100.0)																(100.0)				
腎・尿路結石						2	3	2		3		2	1								13		13	187.60	74.80
(%)						(15.4)	(23.1)	(15.4)		(23.1)		(15.4)	(7.7)								(100.0)				
腎・尿路腫瘍		1		1		1		1	1	1				1					1		7		7	206.60	211.09
(%)		(14.3)		(14.3)		(14.3)		(14.3)	(14.3)	(14.3)				(14.3)					(14.3)		(100.0)				
閉塞性尿路障害				2		1		2		4		1			1					3	14		14	326.36	294.41
(%)				(14.3)		(7.1)		(14.3)		(28.6)		(7.1)		(7.1)						(21.4)	(100.0)				
骨髄腫		1				1				1											3		3	118.33	94.84
(%)		(33.3)				(33.3)				(33.3)											(100.0)				
腎形成不全	2	3	5	1	2	4	3	2	3	5		2	3	1			1	1	1	3	42	1	43	268.74	403.81
(%)	(4.8)	(7.1)	(11.9)	(2.4)	(4.8)	(9.5)	(7.1)	(4.8)	(7.1)	(11.9)		(4.8)	(7.1)	(2.4)			(2.4)	(2.4)	(2.4)	(7.1)	(100.0)				
不明	13	24	24	26	32	30	28	25	17	53	1	43	33	18	17	3	4	7	13	8	419	30	449	227.11	234.40
(%)	(3.1)	(5.7)	(5.7)	(6.2)	(7.6)	(7.2)	(6.7)	(6.0)	(4.1)	(12.6)	(0.2)	(10.3)	(7.9)	(4.3)	(4.1)	(0.7)	(1.0)	(1.7)	(3.1)	(1.9)	(100.0)				
再導入	2	2		5	1	2	2	1	4	4		1		1	3			1	2		31	2	33	211.14	196.09
(%)	(6.5)	(6.5)		(16.1)	(3.2)	(6.5)	(6.5)	(3.2)	(12.9)	(12.9)		(3.2)		(3.2)	(9.7)			(3.2)	(6.5)		(100.0)				
その他	4	9	6	3	6	14	7	6	8	17	1	18	5	2	6	1	1		6	8	128	10	138	290.90	375.65
(%)	(3.1)	(7.0)	(4.7)	(2.3)	(4.7)	(10.9)	(5.5)	(4.7)	(6.3)	(13.3)	(0.8)	(14.1)	(3.9)	(1.6)	(4.7)	(0.8)	(0.8)		(4.7)	(6.3)	(100.0)				
合計	115	230	267	286	271	304	269	268	235	640	9	503	308	177	135	41	52	68	119	118	4,415	328	4,743	230.82	228.91
(%)	(2.6)	(5.2)	(6.0)	(6.5)	(6.1)	(6.9)	(6.1)	(6.1)	(5.3)	(14.5)	(0.2)	(11.4)	(7.0)	(4.0)	(3.1)	(0.9)	(1.2)	(1.5)	(2.7)	(2.7)	(100.0)				
記載なし																									
(%)																									
総計	115	230	267	286	271	304	269	268	235	640	9	503	308	177	135	41	52	68	119	118	4,415	328	4,743	230.82	228.91
(%)	(2.6)	(5.2)	(6.0)	(6.5)	(6.1)	(6.9)	(6.1)	(6.1)	(5.3)	(14.5)	(0.2)	(11.4)	(7.0)	(4.0)	(3.1)	(0.9)	(1.2)	(1.5)	(2.7)	(2.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy
2015年調査