

表 3041 PTH値(pg/mL) 治療方法(新分類)別(透析患者全体) PTH測定法:whole-PTH

治療方法 (新分類)	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
施設血液透析	1,507	1,845	1,688	2,598	2,270	1,796	1,268	41	1,028	703	463	557	217	154	159	207	16,501	629	17,130	108.50	114.02
(%)	(9.1)	(11.2)	(10.2)	(15.7)	(13.8)	(10.9)	(7.7)	(0.2)	(6.2)	(4.3)	(2.8)	(3.4)	(1.3)	(0.9)	(1.0)	(1.3)	(100.0)				
Off-line HDF	31	40	30	45	41	25	20		16	13	4	13	6	4	5	3	296	8	304	105.84	101.28
(%)	(10.5)	(13.5)	(10.1)	(15.2)	(13.9)	(8.4)	(6.8)		(5.4)	(4.4)	(1.4)	(4.4)	(2.0)	(1.4)	(1.7)	(1.0)	(100.0)				
On-line HDF	222	286	324	500	445	341	218	6	211	123	85	97	47	31	41	52	3,029	136	3,165	115.72	129.89
(%)	(7.3)	(9.4)	(10.7)	(16.5)	(14.7)	(11.3)	(7.2)	(0.2)	(7.0)	(4.1)	(2.8)	(3.2)	(1.6)	(1.0)	(1.4)	(1.7)	(100.0)				
Push/Pull HDF				2	2	1	1		1	1							8		8	109.61	49.92
(%)				(25.0)	(25.0)	(12.5)	(12.5)		(12.5)	(12.5)							(100.0)				
ハイオフィльтраーション	3	6	4	6	4	5	2		3		2	2	1			1	39		39	108.51	100.55
(%)	(7.7)	(15.4)	(10.3)	(15.4)	(10.3)	(12.8)	(5.1)		(7.7)		(5.1)	(5.1)	(2.6)			(2.6)	(100.0)				
IHDF	17	21	32	26	37	23	15		10	2	8	6	2	2	4	2	207	1	208	102.76	97.37
(%)	(8.2)	(10.1)	(15.5)	(12.6)	(17.9)	(11.1)	(7.2)		(4.8)	(1.0)	(3.9)	(2.9)	(1.0)	(1.0)	(1.9)	(1.0)	(100.0)				
血液濾過			1														1		1	49.00	
(%)			(100.0)														(100.0)				
血液吸着透析 (リクセル等使用)	19	19	8	21	13	12	11		6	1	2				1	2	115	7	122	83.37	92.77
(%)	(16.5)	(16.5)	(7.0)	(18.3)	(11.3)	(10.4)	(9.6)		(5.2)	(0.9)	(1.7)				(0.9)	(1.7)	(100.0)				
在宅血液透析	3	8	5	12	9	2	3		6	3	2	2				3	58	1	59	124.63	151.17
(%)	(5.2)	(13.8)	(8.6)	(20.7)	(15.5)	(3.4)	(5.2)		(10.3)	(5.2)	(3.4)	(3.4)				(5.2)	(100.0)				
腹膜透析 (CAPD)	29	30	39	69	51	47	37		29	15	15	16	11	7	9	9	413	22	435	130.37	129.26
(%)	(7.0)	(7.3)	(9.4)	(16.7)	(12.3)	(11.4)	(9.0)		(7.0)	(3.6)	(3.6)	(3.9)	(2.7)	(1.7)	(2.2)	(2.2)	(100.0)				
腹膜透析 (APD)	14	13	12	19	21	19	12		12	11	6	8	4		3	4	158	14	172	138.42	161.58
(%)	(8.9)	(8.2)	(7.6)	(12.0)	(13.3)	(12.0)	(7.6)		(7.6)	(7.0)	(3.8)	(5.1)	(2.5)		(1.9)	(2.5)	(100.0)				
腹膜透析 (CCPD)	7	14	5	17	4	6	7	1	7	5	5	4	2		1	3	88	3	91	124.75	116.54
(%)	(8.0)	(15.9)	(5.7)	(19.3)	(4.5)	(6.8)	(8.0)	(1.1)	(8.0)	(5.7)	(5.7)	(4.5)	(2.3)		(1.1)	(3.4)	(100.0)				
合計	1,852	2,282	2,148	3,315	2,897	2,277	1,594	48	1,329	877	592	705	290	198	223	286	20,913	821	21,734	110.08	116.94
(%)	(8.9)	(10.9)	(10.3)	(15.9)	(13.9)	(10.9)	(7.6)	(0.2)	(6.4)	(4.2)	(2.8)	(3.4)	(1.4)	(0.9)	(1.1)	(1.4)	(100.0)				
記載なし																					
総計	1,852	2,282	2,148	3,315	2,897	2,277	1,594	48	1,329	877	592	705	290	198	223	286	20,913	821	21,734	110.08	116.94
(%)	(8.9)	(10.9)	(10.3)	(15.9)	(13.9)	(10.9)	(7.6)	(0.2)	(6.4)	(4.2)	(2.8)	(3.4)	(1.4)	(0.9)	(1.1)	(1.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
 *腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの
 (C)Japanese Society for Dialysis Therapy
 2015年調査