

表 3051 PTH値(pg/mL) 原疾患別（施設血液透析患者 透析回数週3回）PTH測定法-whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	409	529	443	698	636	505	351	12	295	206	128	178	69	36	63	77	4,635	127	4,762	114.18	124.49
(%)	(8.8)	(11.4)	(9.6)	(15.1)	(13.7)	(10.9)	(7.6)	(0.3)	(6.4)	(4.4)	(2.8)	(3.8)	(1.5)	(0.8)	(1.4)	(1.7)	(100.0)				
慢性腎盂腎炎	16	21	28	38	17	26	14	1	10	8	6	7	2	2	1	2	199	5	204	100.82	87.87
(%)	(8.0)	(10.6)	(14.1)	(19.1)	(8.5)	(13.1)	(7.0)	(0.5)	(5.0)	(4.0)	(3.0)	(3.5)	(1.0)	(1.0)	(0.5)	(1.0)	(100.0)				
急速進行性糸球体腎炎	19	30	23	37	18	8	12		8	10	3			4	1		174	10	184	83.86	77.64
(%)	(10.9)	(17.2)	(13.2)	(21.3)	(10.3)	(4.6)	(6.9)		(4.6)	(5.7)	(1.7)			(2.3)	(0.6)		(100.0)				
妊娠腎/妊娠中毒症	5	13	3	7	12	11	7		7	2	1	3	1	1	2	1	76	1	77	120.77	122.40
(%)	(6.6)	(17.1)	(3.9)	(9.2)	(15.8)	(14.5)	(9.2)		(9.2)	(2.6)	(1.3)	(3.9)	(1.3)	(1.3)	(2.6)	(1.3)	(100.0)				
その他の分類不能の腎炎	8	6	6	9	13	8	5		3	2	3	5	1	2			71		71	109.35	90.09
(%)	(11.3)	(8.5)	(8.5)	(12.7)	(18.3)	(11.3)	(7.0)		(4.2)	(2.8)	(4.2)	(7.0)	(1.4)	(2.8)			(100.0)				
多発性嚢胞腎	62	71	62	99	100	64	51	2	39	30	25	27	13	9	5	6	665	12	677	110.39	100.12
(%)	(9.3)	(10.7)	(9.3)	(14.9)	(15.0)	(9.6)	(7.7)	(0.3)	(5.9)	(4.5)	(3.8)	(4.1)	(2.0)	(1.4)	(0.8)	(0.9)	(100.0)				
腎硬化症	129	184	171	262	228	172	145	3	96	68	47	77	15	21	16	7	1,641	66	1,707	106.81	98.72
(%)	(7.9)	(11.2)	(10.4)	(16.0)	(13.9)	(10.5)	(8.8)	(0.2)	(5.9)	(4.1)	(2.9)	(4.7)	(0.9)	(1.3)	(1.0)	(0.4)	(100.0)				
悪性高血圧	13	17	19	24	30	21	9		9	1	9	4	2		2	3	163	5	168	118.50	190.46
(%)	(8.0)	(10.4)	(11.7)	(14.7)	(18.4)	(12.9)	(5.5)		(5.5)	(0.6)	(5.5)	(2.5)	(1.2)		(1.2)	(1.8)	(100.0)				
糖尿病性腎症	588	718	672	1,015	900	682	481	15	409	258	169	167	68	50	44	57	6,293	234	6,527	101.52	97.16
(%)	(9.3)	(11.4)	(10.7)	(16.1)	(14.3)	(10.8)	(7.6)	(0.2)	(6.5)	(4.1)	(2.7)	(2.7)	(1.1)	(0.8)	(0.7)	(0.9)	(100.0)				
SLE腎炎	17	15	10	18	15	12	4		6	3	4	5	2	1		3	115	5	120	108.37	135.16
(%)	(14.8)	(13.0)	(8.7)	(15.7)	(13.0)	(10.4)	(3.5)		(5.2)	(2.6)	(3.5)	(4.3)	(1.7)	(0.9)		(2.6)	(100.0)				
アミロイド腎	3	7	2	7	5	1			4	3	1	1					34		34	89.33	71.24
(%)	(8.8)	(20.6)	(5.9)	(20.6)	(14.7)	(2.9)			(11.8)	(8.8)	(2.9)	(2.9)					(100.0)				
痛風腎	3	3	6	11	7	7	4		8	3	2			2		2	59	4	63	133.41	132.15
(%)	(5.1)	(5.1)	(10.2)	(18.6)	(11.9)	(11.9)	(6.8)		(13.6)	(5.1)	(3.4)			(3.4)		(3.4)	(100.0)				
先天性代謝異常による腎不全		1	1	4	1	2	2				1	1					13		13	105.97	73.05
(%)		(7.7)	(7.7)	(30.8)	(7.7)	(15.4)	(15.4)				(7.7)	(7.7)					(100.0)				
腎・尿路結核	2	2	3	1	2	1			1				1				13		13	93.92	67.13
(%)		(15.4)	(15.4)	(23.1)	(7.7)	(15.4)	(7.7)		(7.7)			(7.7)					(100.0)				
腎・尿路結石	4	2	3	3	6	3	2		2	1	1		1			1	29	2	31	116.04	134.85
(%)	(13.8)	(6.9)	(10.3)	(10.3)	(20.7)	(10.3)	(6.9)		(6.9)	(3.4)	(3.4)		(3.4)			(3.4)	(100.0)				
腎・尿路腫瘍	8	5	6	8	5	6	2		6	1	1	2	1				51	1	52	90.31	71.82
(%)	(15.7)	(9.8)	(11.8)	(15.7)	(9.8)	(11.8)	(3.9)		(11.8)	(2.0)	(2.0)	(3.9)	(2.0)				(100.0)				
閉塞性尿路障害	5	1	6	7	5	2	4		1	2		1	1	1		2	38	2	40	136.73	208.79
(%)	(13.2)	(2.6)	(15.8)	(18.4)	(13.2)	(5.3)	(10.5)		(2.6)	(5.3)		(2.6)	(2.6)	(2.6)		(5.3)	(100.0)				
骨髄腫	2		1	2	2	3	1		2			1					14		14	97.44	60.84
(%)	(14.3)		(7.1)	(14.3)	(14.3)	(21.4)	(7.1)		(14.3)		(7.1)						(100.0)				
腎形成不全	6		6	7	3	6	1		2	3	3		1	1	1	1	41		41	125.74	124.11
(%)	(14.6)		(14.6)	(17.1)	(7.3)	(14.6)	(2.4)		(4.9)	(7.3)	(7.3)		(2.4)	(2.4)	(2.4)	(2.4)	(100.0)				
不明	112	131	118	193	151	166	90	4	73	58	35	45	21	16	14	31	1,258	86	1,344	121.00	140.06
(%)	(8.9)	(10.4)	(9.4)	(15.3)	(12.0)	(13.2)	(7.2)	(0.3)	(5.8)	(4.6)	(2.8)	(3.6)	(1.7)	(1.3)	(1.1)	(2.5)	(100.0)				
再導入	7	6	7	23	16	8	16	2	8	4	6	9	4	1	2	4	123	1	124	150.17	163.51
(%)	(5.7)	(4.9)	(5.7)	(18.7)	(13.0)	(6.5)	(13.0)	(1.6)	(6.5)	(3.3)	(4.9)	(7.3)	(3.3)	(0.8)	(1.6)	(3.3)	(100.0)				
その他	44	43	36	49	53	39	29	2	17	15	6	12	4	5	3	5	362	15	377	106.00	116.62
(%)	(12.2)	(11.9)	(9.9)	(13.5)	(14.6)	(10.8)	(8.0)	(0.6)	(4.7)	(4.1)	(1.7)	(3.3)	(1.1)	(1.4)	(0.8)	(1.4)	(100.0)				
合計	1,460	1,805	1,631	2,524	2,224	1,754	1,231	41	1,006	678	452	545	212	147	155	202	16,067	576	16,643	108.45	112.79
(%)	(9.1)	(11.2)	(10.2)	(15.7)	(13.8)	(10.9)	(7.7)	(0.3)	(6.3)	(4.2)	(2.8)	(3.4)	(1.3)	(0.9)	(1.0)	(1.3)	(100.0)				
記載なし																					
(%)																					
総計	1,460	1,805	1,631	2,524	2,224	1,754	1,231	41	1,006	678	452	545	212	147	155	202	16,067	576	16,643	108.45	112.79
(%)	(9.1)	(11.2)	(10.2)	(15.7)	(13.8)	(10.9)	(7.7)	(0.3)	(6.3)	(4.2)	(2.8)	(3.4)	(1.3)	(0.9)	(1.0)	(1.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査