

表 3056 PTH値(pg/mL) 原疾患別（血液透析導入患者 透析回数週3回）PTH測定法whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	118	120	129	217	201	142	103	2	92	54	39	42	21	14	20	24	1,338	27	1,365	117.36	138.97
(%)	(8.8)	(9.0)	(9.6)	(16.2)	(15.0)	(10.6)	(7.7)	(0.1)	(6.9)	(4.0)	(2.9)	(3.1)	(1.6)	(1.0)	(1.5)	(1.8)	(100.0)				
慢性腎盂腎炎	3	6	4	4	5	3	2		3	1	1	1	2		1	2	38	1	39	131.82	137.98
(%)	(7.9)	(15.8)	(10.5)	(10.5)	(13.2)	(7.9)	(5.3)		(7.9)	(2.6)	(2.6)	(2.6)	(5.3)		(2.6)	(5.3)	(100.0)				
急速進行性糸球体腎炎	4	7		4	3	2	2					1					23		23	62.30	60.28
(%)	(17.4)	(30.4)		(17.4)	(13.0)	(8.7)	(8.7)					(4.3)					(100.0)				
妊娠腎/妊娠中毒症	2	5	3	3	3	1	1		1			1			1		21	3	24	89.05	104.17
(%)	(9.5)	(23.8)	(14.3)	(14.3)	(14.3)	(4.8)	(4.8)		(4.8)			(4.8)			(4.8)		(100.0)				
その他の分類不能の腎炎	3		1	1		4	1			1						1	12		12	133.91	178.65
(%)	(25.0)		(8.3)	(8.3)		(33.3)	(8.3)			(8.3)						(8.3)	(100.0)				
多発性嚢胞腎	9	12	17	16	20	21	16		10	8	9	3	4	3	1	2	151	5	156	123.52	105.24
(%)	(6.0)	(7.9)	(11.3)	(10.6)	(13.2)	(13.9)	(10.6)		(6.6)	(5.3)	(6.0)	(2.0)	(2.6)	(2.0)	(0.7)	(1.3)	(100.0)				
腎硬化症	14	31	35	50	30	28	15		14	7	9	15	1	3	6	4	262	4	266	112.97	109.91
(%)	(5.3)	(11.8)	(13.4)	(19.1)	(11.5)	(10.7)	(5.7)		(5.3)	(2.7)	(3.4)	(5.7)	(0.4)	(1.1)	(2.3)	(1.5)	(100.0)				
悪性高血圧		1	2	6	7	5	3		2								2	2	30	127.49	133.28
(%)		(3.6)	(7.1)	(21.4)	(25.0)	(17.9)	(10.7)		(7.1)								(7.1)	(100.0)			
糖尿病性腎症	70	110	144	182	178	138	79	4	77	41	28	37	16	9	10	8	1,131	51	1,182	104.07	88.53
(%)	(6.2)	(9.7)	(12.7)	(16.1)	(15.7)	(12.2)	(7.0)	(0.4)	(6.8)	(3.6)	(2.5)	(3.3)	(1.4)	(0.8)	(0.9)	(0.7)	(100.0)				
SLE腎炎	7	2	2	3	3	4	1		5	3	1	2	2	1			36	3	39	122.49	100.94
(%)	(19.4)	(5.6)	(5.6)	(8.3)	(8.3)	(11.1)	(2.8)		(13.9)	(8.3)	(2.8)	(5.6)	(5.6)	(2.8)			(100.0)				
アミロイド腎	1	3	2	1		1						1					9		9	70.32	72.42
(%)	(11.1)	(33.3)	(22.2)	(11.1)		(11.1)						(11.1)					(100.0)				
痛風腎	2		1		2	1	2		1	1					1	2	13		13	200.28	203.84
(%)	(15.4)		(7.7)		(15.4)	(7.7)	(15.4)		(7.7)	(7.7)					(7.7)	(15.4)	(100.0)				
先天性代謝異常による腎不全		1	1				1		1								4	1	5	94.22	68.93
(%)		(25.0)	(25.0)				(25.0)		(25.0)								(100.0)				
腎・尿路結核	1			1	1	1	1						1				6		6	112.83	77.82
(%)		(16.7)		(16.7)	(16.7)	(16.7)	(16.7)						(16.7)				(100.0)				
腎・尿路結石				3	2					2							7		7	102.30	62.16
(%)				(42.9)	(28.6)					(28.6)							(100.0)				
腎・尿路腫瘍		1	1		2		1		1								6		6	86.28	48.98
(%)		(16.7)	(16.7)		(33.3)		(16.7)		(16.7)								(100.0)				
閉塞性尿路障害	2			4	2		2			1							11	1	12	80.63	54.67
(%)	(18.2)			(36.4)	(18.2)		(18.2)			(9.1)							(100.0)				
骨髄腫			1			2				1					1		5		5	176.46	146.10
(%)			(20.0)			(40.0)				(20.0)					(20.0)		(100.0)				
腎形成不全	2	1		1	2	1						1				2	10		10	192.59	246.72
(%)	(20.0)	(10.0)		(10.0)	(20.0)	(10.0)						(10.0)				(20.0)	(100.0)				
不明	19	23	23	40	40	23	14		25	10	5	8	6	3	4	6	249	32	281	128.22	167.48
(%)	(7.6)	(9.2)	(9.2)	(16.1)	(16.1)	(9.2)	(5.6)		(10.0)	(4.0)	(2.0)	(3.2)	(2.4)	(1.2)	(1.6)	(2.4)	(100.0)				
再導入	1	4	1	2	5	4	2		1	4	3	2	1	1	1	2	34		34	170.50	136.69
(%)	(2.9)	(11.8)	(2.9)	(5.9)	(14.7)	(11.8)	(5.9)		(2.9)	(11.8)	(8.8)	(5.9)	(2.9)	(2.9)	(2.9)	(5.9)	(100.0)				
その他	7	11	15	20	16	5	6		4	2	2	3	2	3	3	2	101	1	102	130.17	215.68
(%)	(6.9)	(10.9)	(14.9)	(19.8)	(15.8)	(5.0)	(5.9)		(4.0)	(2.0)	(2.0)	(3.0)	(2.0)	(3.0)	(3.0)	(2.0)	(100.0)				
合計	264	339	382	558	522	386	252	6	237	136	97	118	55	37	49	57	3,495	131	3,626	114.72	126.40
(%)	(7.6)	(9.7)	(10.9)	(16.0)	(14.9)	(11.0)	(7.2)	(0.2)	(6.8)	(3.9)	(2.8)	(3.4)	(1.6)	(1.1)	(1.4)	(1.6)	(100.0)				
記載なし																					
(%)																					
総計	264	339	382	558	522	386	252	6	237	136	97	118	55	37	49	57	3,495	131	3,626	114.72	126.40
(%)	(7.6)	(9.7)	(10.9)	(16.0)	(14.9)	(11.0)	(7.2)	(0.2)	(6.8)	(3.9)	(2.8)	(3.4)	(1.6)	(1.1)	(1.4)	(1.6)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査