

表 3060 PTH値(pg/mL) 透析歴(年)別（腹膜透析患者）PTH測定法:whole-PTH

透析歴	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
2年未満	19	12	22	30	21	25	22		24	14	13	14	8	2	6	10	242	6	248	157.55	164.73
(%)	(7.9)	(5.0)	(9.1)	(12.4)	(8.7)	(10.3)	(9.1)		(9.9)	(5.8)	(5.4)	(5.8)	(3.3)	(0.8)	(2.5)	(4.1)	(100.0)				
2年～	16	28	17	27	25	24	16		12	6	7	9	5	2	2	3	199	22	221	114.78	112.91
(%)	(8.0)	(14.1)	(8.5)	(13.6)	(12.6)	(12.1)	(8.0)		(6.0)	(3.0)	(3.5)	(4.5)	(2.5)	(1.0)	(1.0)	(1.5)	(100.0)				
5年～	12	13	13	33	21	16	12	1	10	11	4	3	4	2	5	1	161	9	170	119.03	111.74
(%)	(7.5)	(8.1)	(8.1)	(20.5)	(13.0)	(9.9)	(7.5)	(0.6)	(6.2)	(6.8)	(2.5)	(1.9)	(2.5)	(1.2)	(3.1)	(0.6)	(100.0)				
10年～	1	2	4	11	7	6	5				1	2		1		1	41	2	43	110.01	99.60
(%)	(2.4)	(4.9)	(9.8)	(26.8)	(17.1)	(14.6)	(12.2)				(2.4)	(4.9)		(2.4)		(2.4)	(100.0)				
15年～	2	2		4	2		1				1					1	13		13	122.35	193.49
(%)	(15.4)	(15.4)		(30.8)	(15.4)		(7.7)				(7.7)					(7.7)	(100.0)				
20年～						1			1								2		2	148.40	37.62
(%)						(50.0)			(50.0)								(100.0)				
25年～									1								1		1	162.20	
(%)									(100.0)								(100.0)				
30年～																					
(%)																					
35年～																					
(%)																					
40年～																					
(%)																					
合計	50	57	56	105	76	72	56	1	48	31	26	28	17	7	13	16	659	39	698	131.55	136.03
(%)	(7.6)	(8.6)	(8.5)	(15.9)	(11.5)	(10.9)	(8.5)	(0.2)	(7.3)	(4.7)	(3.9)	(4.2)	(2.6)	(1.1)	(2.0)	(2.4)	(100.0)				
不明																					
(%)																					
記載なし																					
(%)																					
総計	50	57	56	105	76	72	56	1	48	31	26	28	17	7	13	16	659	39	698	131.55	136.03
(%)	(7.6)	(8.6)	(8.5)	(15.9)	(11.5)	(10.9)	(8.5)	(0.2)	(7.3)	(4.7)	(3.9)	(4.2)	(2.6)	(1.1)	(2.0)	(2.4)	(100.0)				
平均	3.40	3.79	3.48	4.71	4.26	4.06	3.68	6.00	3.25	2.94	2.88	2.61	2.65	4.29	3.00	2.94	3.73	3.62	3.73		
標準偏差	3.70	3.35	3.69	4.27	3.99	4.30	4.02		4.89	2.98	3.79	3.08	3.06	4.15	3.21	4.36	3.95	2.55	3.89		

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査