

表 3061 PTH値(pg/mL) 原疾患別（腹膜透析患者）PTH測定法:whole-PTH

原疾患	20未満	20～	35～	50～	75～	100～	125～	150～	151～	180～	210～	240～	300～	350～	400～	500～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	13	24	22	43	28	26	16		17	10	8	11	4	5	9	5	241	11	252	132.31	130.84
(%)	(5.4)	(10.0)	(9.1)	(17.8)	(11.6)	(10.8)	(6.6)		(7.1)	(4.1)	(3.3)	(4.6)	(1.7)	(2.1)	(3.7)	(2.1)	(100.0)				
慢性腎盂腎炎	1	1	3	1		2	1										9		9	68.04	44.18
(%)	(11.1)	(11.1)	(33.3)	(11.1)		(22.2)	(11.1)										(100.0)				
急速進行性糸球体腎炎	1				1				1								3		3	92.10	79.27
(%)	(33.3)				(33.3)				(33.3)								(100.0)				
妊娠腎/妊娠中毒症				3												1	4		4	283.80	447.57
(%)				(75.0)												(25.0)	(100.0)				
その他の分類不能の腎炎				2	1		2										5		5	98.68	37.20
(%)				(40.0)	(20.0)		(40.0)										(100.0)				
多発性嚢胞腎	2		2	3	3	2	1			1	1	1					2		18	162.15	197.63
(%)	(11.1)		(11.1)	(16.7)	(16.7)	(11.1)	(5.6)			(5.6)	(5.6)	(5.6)					(11.1)				
腎硬化症	7	2	6	11	7	8	6		2	2	4	5	4	1	3	2	70	9	79	159.27	174.64
(%)	(10.0)	(2.9)	(8.6)	(15.7)	(10.0)	(11.4)	(8.6)		(2.9)	(2.9)	(5.7)	(7.1)	(5.7)	(1.4)	(4.3)	(2.9)	(100.0)				
悪性高血圧	1	2		1			1			1							6		6	81.47	78.68
(%)	(16.7)	(33.3)		(16.7)			(16.7)			(16.7)							(100.0)				
糖尿病性腎症	19	17	13	31	21	28	18		17	16	7	6	5			1	199	11	210	110.52	79.89
(%)	(9.5)	(8.5)	(6.5)	(15.6)	(10.6)	(14.1)	(9.0)		(8.5)	(8.0)	(3.5)	(3.0)	(2.5)			(0.5)	(100.0)				
SLE腎炎			2			1	2		1		1						7		7	122.43	66.45
(%)			(28.6)			(14.3)	(28.6)		(14.3)		(14.3)						(100.0)				
アミロイド腎												1					1		1	244.00	
(%)												(100.0)					(100.0)				
痛風腎					1	1			1								3		3	125.67	49.60
(%)					(33.3)	(33.3)			(33.3)								(100.0)				
先天性代謝異常による腎不全			1		1												2		2	64.95	37.83
(%)			(50.0)		(50.0)												(100.0)				
腎・尿路結核																					
(%)																					
腎・尿路結石	1	1			1										1		4		4	157.00	226.78
(%)	(25.0)	(25.0)			(25.0)										(25.0)		(100.0)				
腎・尿路腫瘍																					
(%)																					
閉塞性尿路障害																					
(%)																					
骨髄腫					1												1		1	87.00	
(%)					(100.0)												(100.0)				
腎形成不全			1	1	1								1			1	5		5	302.10	404.40
(%)			(20.0)	(20.0)	(20.0)								(20.0)			(20.0)	(100.0)				
不明	4	6	4	8	5	1	7		4	1	4	3	1	1		4	53	7	60	159.62	186.48
(%)	(7.5)	(11.3)	(7.5)	(15.1)	(9.4)	(1.9)	(13.2)		(7.5)	(1.9)	(7.5)	(5.7)	(1.9)	(1.9)		(7.5)	(100.0)				
再導入		1				1											2	1	3	73.50	62.93
(%)		(50.0)				(50.0)											(100.0)				
その他	1	3	2	1	5	2	2	1	5		1	1	2				26		26	126.91	85.20
(%)	(3.8)	(11.5)	(7.7)	(3.8)	(19.2)	(7.7)	(7.7)	(3.8)	(19.2)		(3.8)	(3.8)	(7.7)				(100.0)				
合計	50	57	56	105	76	72	56	1	48	31	26	28	17	7	13	16	659	39	698	131.55	136.03
(%)	(7.6)	(8.6)	(8.5)	(15.9)	(11.5)	(10.9)	(8.5)	(0.2)	(7.3)	(4.7)	(3.9)	(4.2)	(2.6)	(1.1)	(2.0)	(2.4)	(100.0)				
記載なし																					
(%)																					
総計	50	57	56	105	76	72	56	1	48	31	26	28	17	7	13	16	659	39	698	131.55	136.03
(%)	(7.6)	(8.6)	(8.5)	(15.9)	(11.5)	(10.9)	(8.5)	(0.2)	(7.3)	(4.7)	(3.9)	(4.2)	(2.6)	(1.1)	(2.0)	(2.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査