

表 3064 透析前ヘモグロビン濃度(g/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
施設血液透析	219	387	960	3,512	11,682	36,375	75,469	63,774	24,012	6,047	2,155	224,592	24,133	248,725	10.72	1.29
(%)	(0.1)	(0.2)	(0.4)	(1.6)	(5.2)	(16.2)	(33.6)	(28.4)	(10.7)	(2.7)	(1.0)	(100.0)				
Off-line HDF	4	8	20	80	269	803	1,521	1,304	554	157	52	4,772	560	5,332	10.73	1.32
(%)	(0.1)	(0.2)	(0.4)	(1.7)	(5.6)	(16.8)	(31.9)	(27.3)	(11.6)	(3.3)	(1.1)	(100.0)				
On-line HDF	27	32	93	394	1,418	5,537	13,727	12,673	4,860	1,348	584	40,693	3,834	44,527	10.91	1.24
(%)	(0.1)	(0.1)	(0.2)	(1.0)	(3.5)	(13.6)	(33.7)	(31.1)	(11.9)	(3.3)	(1.4)	(100.0)				
Push/Pull HDF					3	21	43	24	11	3	2	107	3	110	10.81	1.16
(%)					(2.8)	(19.6)	(40.2)	(22.4)	(10.3)	(2.8)	(1.9)	(100.0)				
バイオフィльтраーション		1		4	6	39	79	78	33	7	3	250	17	267	10.89	1.26
(%)		(0.4)		(1.6)	(2.4)	(15.6)	(31.6)	(31.2)	(13.2)	(2.8)	(1.2)	(100.0)				
IHDF	3	2	5	35	176	658	1,162	932	323	67	19	3,382	158	3,540	10.66	1.18
(%)	(0.1)	(0.1)	(0.1)	(1.0)	(5.2)	(19.5)	(34.4)	(27.6)	(9.6)	(2.0)	(0.6)	(100.0)				
血液濾過		1		2	2		4	3				12	4	16	9.54	1.91
(%)		(8.3)		(16.7)	(16.7)		(33.3)	(25.0)				(100.0)				
血液吸着透析 (リクセル等使用)		1	5	21	66	239	433	385	130	23	10	1,313	63	1,376	10.67	1.20
(%)		(0.1)	(0.4)	(1.6)	(5.0)	(18.2)	(33.0)	(29.3)	(9.9)	(1.8)	(0.8)	(100.0)				
在宅血液透析		3	2		14	45	102	126	81	42	23	438	117	555	11.43	1.56
(%)		(0.7)	(0.5)		(3.2)	(10.3)	(23.3)	(28.8)	(18.5)	(9.6)	(5.3)	(100.0)				
腹膜透析 (CAPD)	1	3	20	62	193	516	919	910	442	122	47	3,235	1,667	4,902	10.83	1.41
(%)	(0.0)	(0.1)	(0.6)	(1.9)	(6.0)	(16.0)	(28.4)	(28.1)	(13.7)	(3.8)	(1.5)	(100.0)				
腹膜透析 (APD)	1	1	10	26	131	304	578	524	263	81	25	1,944	831	2,775	10.82	1.37
(%)	(0.1)	(0.1)	(0.5)	(1.3)	(6.7)	(15.6)	(29.7)	(27.0)	(13.5)	(4.2)	(1.3)	(100.0)				
腹膜透析 (CCPD)		1	9	9	37	134	206	245	97	27	11	776	316	1,092	10.83	1.37
(%)		(0.1)	(1.2)	(1.2)	(4.8)	(17.3)	(26.5)	(31.6)	(12.5)	(3.5)	(1.4)	(100.0)				
合計	255	440	1,124	4,145	13,997	44,671	94,243	80,978	30,806	7,924	2,931	281,514	31,703	313,217	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				
記載なし (%)																
総計	255	440	1,124	4,145	13,997	44,671	94,243	80,978	30,806	7,924	2,931	281,514	31,703	313,217	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査