

表 3065 透析前ヘモグロビン濃度(g/dL) 併用の状況別（透析患者全体）

併用の状況	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	253	435	1,083	4,040	13,617	43,674	92,468	79,240	29,962	7,689	2,841	275,302	28,861	304,163	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(4.9)	(15.9)	(33.6)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				
PD単独	1	4	32	81	281	769	1,350	1,276	613	162	67	4,636	2,468	7,104	10.79	1.40
(%)	(0.0)	(0.1)	(0.7)	(1.7)	(6.1)	(16.6)	(29.1)	(27.5)	(13.2)	(3.5)	(1.4)	(100.0)				
HD(F)単独.PDカテ あり			2	6	13	26	46	38	26	3	1	161	21	182	10.63	1.45
(%)			(1.2)	(3.7)	(8.1)	(16.1)	(28.6)	(23.6)	(16.1)	(1.9)	(0.6)	(100.0)				
PD.HD併用週1 HD(F)	1	1	6	14	70	168	305	356	166	62	14	1,163	307	1,470	10.94	1.37
(%)	(0.1)	(0.1)	(0.5)	(1.2)	(6.0)	(14.4)	(26.2)	(30.6)	(14.3)	(5.3)	(1.2)	(100.0)				
PD.HD併用週2 HD(F)				2	8	17	37	40	25	7	7	143	29	172	11.21	1.51
(%)				(1.4)	(5.6)	(11.9)	(25.9)	(28.0)	(17.5)	(4.9)	(4.9)	(100.0)				
PD.HD併用週3 HD(F)				2		6	9	7	2			26	3	29	10.35	1.19
(%)				(7.7)		(23.1)	(34.6)	(26.9)	(7.7)			(100.0)				
PD.HD併用週4 HD(F)																
(%)																
PD.HD併用その他			1		6	7	22	16	8	1		61	10	71	10.66	1.27
(%)			(1.6)		(9.8)	(11.5)	(36.1)	(26.2)	(13.1)	(1.6)		(100.0)				
合計	255	440	1,124	4,145	13,995	44,667	94,237	80,973	30,802	7,924	2,930	281,492	31,699	313,191	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				
不明					2	4	6	5	4		1	22	3	25	10.86	1.39
(%)					(9.1)	(18.2)	(27.3)	(22.7)	(18.2)		(4.5)	(100.0)				
記載なし													1	1		
(%)																
総計	255	440	1,124	4,145	13,997	44,671	94,243	80,978	30,806	7,924	2,931	281,514	31,703	313,217	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査