

表 3069 透析前ヘモグロビン濃度(g/dL) 原疾患別(透析患者全体)

原疾患	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	85	122	285	1,050	3,770	12,961	28,264	24,834	9,389	2,357	855	83,972	9,375	93,347	10.79	1.26
(%)	(0.1)	(0.1)	(0.3)	(1.3)	(4.5)	(15.4)	(33.7)	(29.6)	(11.2)	(2.8)	(1.0)	(100.0)				
慢性腎盂腎炎	1	4	8	51	146	458	891	740	283	67	26	2,675	260	2,935	10.70	1.28
(%)	(0.0)	(0.1)	(0.3)	(1.9)	(5.5)	(17.1)	(33.3)	(27.7)	(10.6)	(2.5)	(1.0)	(100.0)				
急速進行性糸球体腎炎	2	2	12	31	122	333	737	634	247	74	21	2,215	263	2,478	10.76	1.29
(%)	(0.1)	(0.1)	(0.5)	(1.4)	(5.5)	(15.0)	(33.3)	(28.6)	(11.2)	(3.3)	(0.9)	(100.0)				
妊娠腎/妊娠中毒症		3	5	16	77	238	481	428	152	25	12	1,437	108	1,545	10.72	1.21
(%)		(0.2)	(0.3)	(1.1)	(5.4)	(16.6)	(33.5)	(29.8)	(10.6)	(1.7)	(0.8)	(100.0)				
その他の分類不能の腎炎	1	2	10	22	61	180	434	379	136	36	14	1,275	130	1,405	10.75	1.31
(%)	(0.1)	(0.2)	(0.8)	(1.7)	(4.8)	(14.1)	(34.0)	(29.7)	(10.7)	(2.8)	(1.1)	(100.0)				
多発性嚢胞腎	6	17	17	103	365	1,350	3,339	3,145	1,326	354	199	10,221	1,035	11,256	10.96	1.30
(%)	(0.1)	(0.2)	(0.2)	(1.0)	(3.6)	(13.2)	(32.7)	(30.8)	(13.0)	(3.5)	(1.9)	(100.0)				
腎硬化症	30	36	109	420	1,387	4,413	9,210	7,656	2,699	644	218	26,822	2,983	29,805	10.70	1.26
(%)	(0.1)	(0.1)	(0.4)	(1.6)	(5.2)	(16.5)	(34.3)	(28.5)	(10.1)	(2.4)	(0.8)	(100.0)				
悪性高血圧	1	1	8	25	115	373	718	723	275	69	27	2,335	283	2,618	10.82	1.25
(%)	(0.0)	(0.0)	(0.3)	(1.1)	(4.9)	(16.0)	(30.7)	(31.0)	(11.8)	(3.0)	(1.2)	(100.0)				
糖尿病性腎症	107	184	458	1,638	5,554	17,561	36,154	30,492	11,811	3,183	1,166	108,308	11,970	120,278	10.74	1.30
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.1)	(16.2)	(33.4)	(28.2)	(10.9)	(2.9)	(1.1)	(100.0)				
SLE腎炎	2	3	8	28	113	353	683	541	194	51	16	1,992	235	2,227	10.67	1.26
(%)	(0.1)	(0.2)	(0.4)	(1.4)	(5.7)	(17.7)	(34.3)	(27.2)	(9.7)	(2.6)	(0.8)	(100.0)				
アミロイド腎			4	10	27	72	111	116	50	9	3	402	60	462	10.67	1.35
(%)			(1.0)	(2.5)	(6.7)	(17.9)	(27.6)	(28.9)	(12.4)	(2.2)	(0.7)	(100.0)				
痛風腎	1		5	14	48	148	314	305	105	27	13	980	95	1,075	10.80	1.30
(%)	(0.1)		(0.5)	(1.4)	(4.9)	(15.1)	(32.0)	(31.1)	(10.7)	(2.8)	(1.3)	(100.0)				
先天性代謝異常による腎不全		1	1	5	13	38	71	70	25	8	4	236	38	274	10.75	1.41
(%)		(0.4)	(0.4)	(2.1)	(5.5)	(16.1)	(30.1)	(29.7)	(10.6)	(3.4)	(1.7)	(100.0)				
腎・尿路結核	1	1	2	3	12	37	49	55	14		1	175	20	195	10.44	1.34
(%)	(0.6)	(0.6)	(1.1)	(1.7)	(6.9)	(21.1)	(28.0)	(31.4)	(8.0)		(0.6)	(100.0)				
腎・尿路結石	1		3	8	30	78	172	151	65	12	5	525	56	581	10.74	1.30
(%)	(0.2)		(0.6)	(1.5)	(5.7)	(14.9)	(32.8)	(28.8)	(12.4)	(2.3)	(1.0)	(100.0)				
腎・尿路腫瘍			5	30	55	149	278	208	70	11	12	818	92	910	10.50	1.34
(%)			(0.6)	(3.7)	(6.7)	(18.2)	(34.0)	(25.4)	(8.6)	(1.3)	(1.5)	(100.0)				
閉塞性尿路障害		2	4	8	36	95	216	176	84	26	5	652	78	730	10.80	1.32
(%)		(0.3)	(0.6)	(1.2)	(5.5)	(14.6)	(33.1)	(27.0)	(12.9)	(4.0)	(0.8)	(100.0)				
骨髄腫		1	6	22	35	49	68	44	26	3		254	37	291	10.00	1.56
(%)		(0.4)	(2.4)	(8.7)	(13.8)	(19.3)	(26.8)	(17.3)	(10.2)	(1.2)		(100.0)				
腎形成不全			4	6	24	80	191	186	73	23	10	597	63	660	10.91	1.31
(%)			(0.7)	(1.0)	(4.0)	(13.4)	(32.0)	(31.2)	(12.2)	(3.9)	(1.7)	(100.0)				
不明	12	40	127	474	1,516	4,255	8,855	7,541	2,780	672	232	26,504	3,393	29,897	10.70	1.28
(%)	(0.0)	(0.2)	(0.5)	(1.8)	(5.7)	(16.1)	(33.4)	(28.5)	(10.5)	(2.5)	(0.9)	(100.0)				
再導入	1	6	4	24	80	297	633	566	235	62	22	1,930	253	2,183	10.84	1.27
(%)	(0.1)	(0.3)	(0.2)	(1.2)	(4.1)	(15.4)	(32.8)	(29.3)	(12.2)	(3.2)	(1.1)	(100.0)				
その他	4	15	39	155	411	1,150	2,369	1,986	765	210	70	7,174	870	8,044	10.70	1.33
(%)	(0.1)	(0.2)	(0.5)	(2.2)	(5.7)	(16.0)	(33.0)	(27.7)	(10.7)	(2.9)	(1.0)	(100.0)				
合計	255	440	1,124	4,143	13,997	44,668	94,238	80,976	30,804	7,923	2,931	281,499	31,697	313,196	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				
記載なし				2		3	5	2	2	1		15	6	21	10.43	1.55
(%)				(13.3)		(20.0)	(33.3)	(13.3)	(13.3)	(6.7)		(100.0)				
総計	255	440	1,124	4,145	13,997	44,671	94,243	80,978	30,806	7,924	2,931	281,514	31,703	313,217	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査