

表 3070 透析前ヘモグロビン濃度(g/dL) 医療機関別（透析患者全体）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学		3	9	35	96	169	224	188	81	25	5	835	373	1,208	10.36	1.49
(%)		(0.4)	(1.1)	(4.2)	(11.5)	(20.2)	(26.8)	(22.5)	(9.7)	(3.0)	(0.6)	(100.0)				
私立大学	2	5	27	74	190	398	738	710	251	68	22	2,485	1,079	3,564	10.61	1.43
(%)	(0.1)	(0.2)	(1.1)	(3.0)	(7.6)	(16.0)	(29.7)	(28.6)	(10.1)	(2.7)	(0.9)	(100.0)				
国立		1	12	35	78	123	175	150	79	17	11	681	243	924	10.41	1.63
(%)		(0.1)	(1.8)	(5.1)	(11.5)	(18.1)	(25.7)	(22.0)	(11.6)	(2.5)	(1.6)	(100.0)				
県市町村立	11	33	122	398	1,069	2,679	5,013	4,479	1,879	543	227	16,453	3,065	19,518	10.70	1.41
(%)	(0.1)	(0.2)	(0.7)	(2.4)	(6.5)	(16.3)	(30.5)	(27.2)	(11.4)	(3.3)	(1.4)	(100.0)				
社会保険	1	3	20	79	207	554	1,097	920	389	112	38	3,420	479	3,899	10.71	1.36
(%)	(0.0)	(0.1)	(0.6)	(2.3)	(6.1)	(16.2)	(32.1)	(26.9)	(11.4)	(3.3)	(1.1)	(100.0)				
厚生連	15	74	62	153	433	1,166	2,483	2,210	995	307	101	7,999	1,002	9,001	10.74	1.47
(%)	(0.2)	(0.9)	(0.8)	(1.9)	(5.4)	(14.6)	(31.0)	(27.6)	(12.4)	(3.8)	(1.3)	(100.0)				
その他公的	12	25	49	184	533	1,308	2,555	2,358	1,106	303	122	8,555	2,369	10,924	10.76	1.43
(%)	(0.1)	(0.3)	(0.6)	(2.2)	(6.2)	(15.3)	(29.9)	(27.6)	(12.9)	(3.5)	(1.4)	(100.0)				
私立総合	6	6	42	97	317	942	2,011	1,894	768	226	75	6,384	1,064	7,448	10.82	1.33
(%)	(0.1)	(0.1)	(0.7)	(1.5)	(5.0)	(14.8)	(31.5)	(29.7)	(12.0)	(3.5)	(1.2)	(100.0)				
私立	40	94	395	1,521	4,677	13,439	25,655	21,972	9,137	2,470	955	80,355	9,710	90,065	10.72	1.33
(%)	(0.0)	(0.1)	(0.5)	(1.9)	(5.8)	(16.7)	(31.9)	(27.3)	(11.4)	(3.1)	(1.2)	(100.0)				
私立診療所	168	196	386	1,569	6,397	23,893	54,292	46,097	16,121	3,853	1,375	154,347	12,319	166,666	10.78	1.21
(%)	(0.1)	(0.1)	(0.3)	(1.0)	(4.1)	(15.5)	(35.2)	(29.9)	(10.4)	(2.5)	(0.9)	(100.0)				
総計	255	440	1,124	4,145	13,997	44,671	94,243	80,978	30,806	7,924	2,931	281,514	31,703	313,217	10.75	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.0)	(15.9)	(33.5)	(28.8)	(10.9)	(2.8)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査