

表 3082 透析前ヘモグロビン濃度(g/dL) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学		2	6	22	59	92	103	76	23	11	3	397	11	408	10.07	1.55
(%)		(0.5)	(1.5)	(5.5)	(14.9)	(23.2)	(25.9)	(19.1)	(5.8)	(2.8)	(0.8)	(100.0)				
私立大学	1	5	21	57	126	231	421	398	136	39	12	1,447	49	1,496	10.51	1.47
(%)	(0.1)	(0.3)	(1.5)	(3.9)	(8.7)	(16.0)	(29.1)	(27.5)	(9.4)	(2.7)	(0.8)	(100.0)				
国立			9	24	58	76	116	85	51	12	4	435	80	515	10.33	1.62
(%)			(2.1)	(5.5)	(13.3)	(17.5)	(26.7)	(19.5)	(11.7)	(2.8)	(0.9)	(100.0)				
県市町村立	10	30	85	309	828	2,021	3,837	3,260	1,376	390	156	12,302	668	12,970	10.67	1.41
(%)	(0.1)	(0.2)	(0.7)	(2.5)	(6.7)	(16.4)	(31.2)	(26.5)	(11.2)	(3.2)	(1.3)	(100.0)				
社会保険	1	3	13	60	153	431	860	699	293	90	31	2,634	41	2,675	10.72	1.36
(%)	(0.0)	(0.1)	(0.5)	(2.3)	(5.8)	(16.4)	(32.6)	(26.5)	(11.1)	(3.4)	(1.2)	(100.0)				
厚生連	13	69	44	120	332	899	1,921	1,710	764	227	66	6,165	258	6,423	10.72	1.48
(%)	(0.2)	(1.1)	(0.7)	(1.9)	(5.4)	(14.6)	(31.2)	(27.7)	(12.4)	(3.7)	(1.1)	(100.0)				
その他公的	9	15	37	142	406	981	1,965	1,793	823	223	98	6,492	424	6,916	10.76	1.43
(%)	(0.1)	(0.2)	(0.6)	(2.2)	(6.3)	(15.1)	(30.3)	(27.6)	(12.7)	(3.4)	(1.5)	(100.0)				
私立総合	6	4	35	75	213	662	1,422	1,359	561	154	52	4,543	246	4,789	10.82	1.35
(%)	(0.1)	(0.1)	(0.8)	(1.7)	(4.7)	(14.6)	(31.3)	(29.9)	(12.3)	(3.4)	(1.1)	(100.0)				
私立	38	74	338	1,245	3,756	10,894	20,505	17,467	7,098	1,867	706	63,988	2,609	66,597	10.70	1.33
(%)	(0.1)	(0.1)	(0.5)	(1.9)	(5.9)	(17.0)	(32.0)	(27.3)	(11.1)	(2.9)	(1.1)	(100.0)				
私立診療所	136	166	308	1,227	5,085	18,601	41,794	34,855	12,014	2,788	936	117,910	3,153	121,063	10.75	1.20
(%)	(0.1)	(0.1)	(0.3)	(1.0)	(4.3)	(15.8)	(35.4)	(29.6)	(10.2)	(2.4)	(0.8)	(100.0)				
総計	214	368	896	3,281	11,016	34,888	72,944	61,702	23,139	5,801	2,064	216,313	7,539	223,852	10.73	1.28
(%)	(0.1)	(0.2)	(0.4)	(1.5)	(5.1)	(16.1)	(33.7)	(28.5)	(10.7)	(2.7)	(1.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査