

表 3094 透析前ヘモグロビン濃度(g/dL) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学				6	10	11	5	11	3	1		47	2	49	9.86	1.55
(%)				(12.8)	(21.3)	(23.4)	(10.6)	(23.4)	(6.4)	(2.1)		(100.0)				
私立大学			1	1	5	10	68	65	17	4	2	173	2	175	10.97	1.09
(%)			(0.6)	(0.6)	(2.9)	(5.8)	(39.3)	(37.6)	(9.8)	(2.3)	(1.2)	(100.0)				
国立			1	3	3	11	13	11	2	4	2	50	1	51	10.49	1.86
(%)			(2.0)	(6.0)	(6.0)	(22.0)	(26.0)	(22.0)	(4.0)	(8.0)	(4.0)	(100.0)				
県市町村立	1		13	34	95	290	580	627	260	81	36	2,017	98	2,115	10.89	1.36
(%)	(0.0)		(0.6)	(1.7)	(4.7)	(14.4)	(28.8)	(31.1)	(12.9)	(4.0)	(1.8)	(100.0)				
社会保険			2	5	21	50	131	128	51	10	1	399	14	413	10.81	1.18
(%)			(0.5)	(1.3)	(5.3)	(12.5)	(32.8)	(32.1)	(12.8)	(2.5)	(0.3)	(100.0)				
厚生連		1	4	9	40	135	313	283	129	45	20	979	43	1,022	10.96	1.33
(%)		(0.1)	(0.4)	(0.9)	(4.1)	(13.8)	(32.0)	(28.9)	(13.2)	(4.6)	(2.0)	(100.0)				
その他公的	2			7	33	108	228	214	124	31	9	756	98	854	10.97	1.32
(%)	(0.3)			(0.9)	(4.4)	(14.3)	(30.2)	(28.3)	(16.4)	(4.1)	(1.2)	(100.0)				
私立総合		2	5	11	70	193	394	354	130	38	15	1,212	30	1,242	10.77	1.27
(%)		(0.2)	(0.4)	(0.9)	(5.8)	(15.9)	(32.5)	(29.2)	(10.7)	(3.1)	(1.2)	(100.0)				
私立	1	14	35	156	525	1,716	3,764	3,400	1,472	434	182	11,699	555	12,254	10.87	1.30
(%)	(0.0)	(0.1)	(0.3)	(1.3)	(4.5)	(14.7)	(32.2)	(29.1)	(12.6)	(3.7)	(1.6)	(100.0)				
私立診療所	30	24	53	260	1,013	4,340	10,670	9,633	3,462	885	374	30,744	662	31,406	10.88	1.20
(%)	(0.1)	(0.1)	(0.2)	(0.8)	(3.3)	(14.1)	(34.7)	(31.3)	(11.3)	(2.9)	(1.2)	(100.0)				
総計	34	41	114	492	1,815	6,864	16,166	14,726	5,650	1,533	641	48,076	1,505	49,581	10.88	1.24
(%)	(0.1)	(0.1)	(0.2)	(1.0)	(3.8)	(14.3)	(33.6)	(30.6)	(11.8)	(3.2)	(1.3)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査