

表 3106 ヘモグロビン濃度(g/dL) 医療機関別（腹膜透析患者）

医療機関	5未満	5～	6～	7～	8～	9～	10～	11～	12～	13～	14～	合計	記載なし	総計	平均	標準偏差
国公立大学			2	4	22	55	108	97	55	13	2	358	305	663	10.83	1.27
(%)			(0.6)	(1.1)	(6.1)	(15.4)	(30.2)	(27.1)	(15.4)	(3.6)	(0.6)	(100.0)				
私立大学			2	7	30	120	202	193	83	20	5	662	444	1,106	10.83	1.28
(%)			(0.3)	(1.1)	(4.5)	(18.1)	(30.5)	(29.2)	(12.5)	(3.0)	(0.8)	(100.0)				
国立			2	2	10	28	40	50	21	1	5	159	63	222	10.81	1.49
(%)			(1.3)	(1.3)	(6.3)	(17.6)	(25.2)	(31.4)	(13.2)	(0.6)	(3.1)	(100.0)				
県市町村立			10	19	70	182	328	326	134	41	17	1,127	446	1,573	10.79	1.40
(%)			(0.9)	(1.7)	(6.2)	(16.1)	(29.1)	(28.9)	(11.9)	(3.6)	(1.5)	(100.0)				
社会保険			3	7	17	34	45	47	23	8	4	188	334	522	10.60	1.57
(%)			(1.6)	(3.7)	(9.0)	(18.1)	(23.9)	(25.0)	(12.2)	(4.3)	(2.1)	(100.0)				
厚生連			5	10	37	65	122	113	47	22	6	427	133	560	10.72	1.47
(%)			(1.2)	(2.3)	(8.7)	(15.2)	(28.6)	(26.5)	(11.0)	(5.2)	(1.4)	(100.0)				
その他公的	1	4	8	14	51	150	269	271	127	33	13	941	418	1,359	10.80	1.42
(%)	(0.1)	(0.4)	(0.9)	(1.5)	(5.4)	(15.9)	(28.6)	(28.8)	(13.5)	(3.5)	(1.4)	(100.0)				
私立総合			1	4	12	32	88	101	48	21	4	311	52	363	11.13	1.33
(%)			(0.3)	(1.3)	(3.9)	(10.3)	(28.3)	(32.5)	(15.4)	(6.8)	(1.3)	(100.0)				
私立		1	4	23	92	197	344	301	188	52	18	1,220	479	1,699	10.82	1.44
(%)		(0.1)	(0.3)	(1.9)	(7.5)	(16.1)	(28.2)	(24.7)	(15.4)	(4.3)	(1.5)	(100.0)				
私立診療所	1		2	7	20	91	157	180	76	19	9	562	140	702	10.92	1.30
(%)	(0.2)		(0.4)	(1.2)	(3.6)	(16.2)	(27.9)	(32.0)	(13.5)	(3.4)	(1.6)	(100.0)				
総計	2	5	39	97	361	954	1,703	1,679	802	230	83	5,955	2,814	8,769	10.83	1.39
(%)	(0.0)	(0.1)	(0.7)	(1.6)	(6.1)	(16.0)	(28.6)	(28.2)	(13.5)	(3.9)	(1.4)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査