

表 3115 総コレステロール濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
施設血液透析	7,130	20,574	37,237	43,083	35,120	22,182	11,275	5,081	2,087	1,365	185,134	63,591	248,725	155.13	35.60
(%)	(3.9)	(11.1)	(20.1)	(23.3)	(19.0)	(12.0)	(6.1)	(2.7)	(1.1)	(0.7)	(100.0)				
Off-line HDF	185	376	715	863	736	472	232	109	60	30	3,778	1,554	5,332	156.04	36.49
(%)	(4.9)	(10.0)	(18.9)	(22.8)	(19.5)	(12.5)	(6.1)	(2.9)	(1.6)	(0.8)	(100.0)				
On-line HDF	909	3,004	6,230	7,795	6,807	4,644	2,524	1,182	437	276	33,808	10,719	44,527	159.75	35.65
(%)	(2.7)	(8.9)	(18.4)	(23.1)	(20.1)	(13.7)	(7.5)	(3.5)	(1.3)	(0.8)	(100.0)				
Push/Pull HDF	1	6	12	31	17	14	7	4	1	2	95	15	110	165.31	35.16
(%)	(1.1)	(6.3)	(12.6)	(32.6)	(17.9)	(14.7)	(7.4)	(4.2)	(1.1)	(2.1)	(100.0)				
ハイオフィルトレーション	8	14	46	44	35	31	15	9	2	1	205	62	267	158.25	35.85
(%)	(3.9)	(6.8)	(22.4)	(21.5)	(17.1)	(15.1)	(7.3)	(4.4)	(1.0)	(0.5)	(100.0)				
IHDF	91	292	569	711	522	346	196	88	32	17	2,864	676	3,540	156.56	35.03
(%)	(3.2)	(10.2)	(19.9)	(24.8)	(18.2)	(12.1)	(6.8)	(3.1)	(1.1)	(0.6)	(100.0)				
血液濾過	2		2	1	4	1					10	6	16	140.60	43.17
(%)	(20.0)		(20.0)	(10.0)	(40.0)	(10.0)					(100.0)				
血液吸着透析 (リクセル等使用)	30	107	199	254	225	169	90	35	16	6	1,131	245	1,376	160.18	35.10
(%)	(2.7)	(9.5)	(17.6)	(22.5)	(19.9)	(14.9)	(8.0)	(3.1)	(1.4)	(0.5)	(100.0)				
在宅血液透析	1	8	35	58	87	80	44	24	13	12	362	193	555	181.29	37.34
(%)	(0.3)	(2.2)	(9.7)	(16.0)	(24.0)	(22.1)	(12.2)	(6.6)	(3.6)	(3.3)	(100.0)				
腹膜透析 (CAPD)	27	114	226	428	479	467	345	201	101	85	2,473	2,429	4,902	180.16	40.66
(%)	(1.1)	(4.6)	(9.1)	(17.3)	(19.4)	(18.9)	(14.0)	(8.1)	(4.1)	(3.4)	(100.0)				
腹膜透析 (APD)	18	65	184	260	356	246	163	103	51	43	1,489	1,286	2,775	175.52	40.82
(%)	(1.2)	(4.4)	(12.4)	(17.5)	(23.9)	(16.5)	(10.9)	(6.9)	(3.4)	(2.9)	(100.0)				
腹膜透析 (CCPD)	7	26	76	139	127	98	69	34	21	23	620	472	1,092	174.31	40.35
(%)	(1.1)	(4.2)	(12.3)	(22.4)	(20.5)	(15.8)	(11.1)	(5.5)	(3.4)	(3.7)	(100.0)				
合計	8,409	24,586	45,531	53,667	44,515	28,750	14,960	6,870	2,821	1,860	231,969	81,248	313,217	156.35	35.90
(%)	(3.6)	(10.6)	(19.6)	(23.1)	(19.2)	(12.4)	(6.4)	(3.0)	(1.2)	(0.8)	(100.0)				
記載なし															
(%)															
総計	8,409	24,586	45,531	53,667	44,515	28,750	14,960	6,870	2,821	1,860	231,969	81,248	313,217	156.35	35.90
(%)	(3.6)	(10.6)	(19.6)	(23.1)	(19.2)	(12.4)	(6.4)	(3.0)	(1.2)	(0.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査