

表 3120 総コレステロール濃度(mg/dL) 原疾患別 (透析患者全体)

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,872	5,875	12,356	15,947	14,400	9,876	5,272	2,469	955	604	69,626	23,721	93,347	160.58	35.65
(%)	(2.7)	(8.4)	(17.7)	(22.9)	(20.7)	(14.2)	(7.6)	(3.5)	(1.4)	(0.9)	(100.0)				
慢性腎盂腎炎	52	183	337	496	465	334	193	95	49	23	2,227	708	2,935	164.06	36.73
(%)	(2.3)	(8.2)	(15.1)	(22.3)	(20.9)	(15.0)	(8.7)	(4.3)	(2.2)	(1.0)	(100.0)				
急速進行性糸球体腎炎	30	97	275	357	364	275	196	123	50	54	1,821	657	2,478	171.95	41.05
(%)	(1.6)	(5.3)	(15.1)	(19.6)	(20.0)	(15.1)	(10.8)	(6.8)	(2.7)	(3.0)	(100.0)				
妊娠腎/妊娠中毒症	10	41	139	217	279	222	158	75	23	19	1,183	362	1,545	174.45	35.27
(%)	(0.8)	(3.5)	(11.7)	(18.3)	(23.6)	(18.8)	(13.4)	(6.3)	(1.9)	(1.6)	(100.0)				
その他の分類不能の腎炎	36	93	179	217	212	135	92	47	16	12	1,039	366	1,405	161.52	38.52
(%)	(3.5)	(9.0)	(17.2)	(20.9)	(20.4)	(13.0)	(8.9)	(4.5)	(1.5)	(1.2)	(100.0)				
多発性嚢胞腎	207	651	1,425	2,052	1,823	1,200	664	252	116	65	8,455	2,801	11,256	161.03	34.57
(%)	(2.4)	(7.7)	(16.9)	(24.3)	(21.6)	(14.2)	(7.9)	(3.0)	(1.4)	(0.8)	(100.0)				
腎硬化症	660	2,266	4,386	5,438	4,371	2,736	1,385	633	243	147	22,265	7,540	29,805	156.41	34.68
(%)	(3.0)	(10.2)	(19.7)	(24.4)	(19.6)	(12.3)	(6.2)	(2.8)	(1.1)	(0.7)	(100.0)				
悪性高血圧	34	148	355	436	421	261	136	65	29	26	1,911	707	2,618	161.70	36.32
(%)	(1.8)	(7.7)	(18.6)	(22.8)	(22.0)	(13.7)	(7.1)	(3.4)	(1.5)	(1.4)	(100.0)				
糖尿病性腎症	4,438	11,977	19,871	20,765	15,457	9,121	4,440	1,960	773	557	89,359	30,919	120,278	150.52	35.08
(%)	(5.0)	(13.4)	(22.2)	(23.2)	(17.3)	(10.2)	(5.0)	(2.2)	(0.9)	(0.6)	(100.0)				
SLE腎炎	23	110	199	331	348	284	176	85	61	29	1,646	581	2,227	171.42	38.68
(%)	(1.4)	(6.7)	(12.1)	(20.1)	(21.1)	(17.3)	(10.7)	(5.2)	(3.7)	(1.8)	(100.0)				
アミロイド腎	11	34	47	57	60	47	32	11	9	5	313	149	462	163.85	40.90
(%)	(3.5)	(10.9)	(15.0)	(18.2)	(19.2)	(15.0)	(10.2)	(3.5)	(2.9)	(1.6)	(100.0)				
痛風腎	29	79	166	188	162	101	54	25	11	3	818	257	1,075	156.75	34.36
(%)	(3.5)	(9.7)	(20.3)	(23.0)	(19.8)	(12.3)	(6.6)	(3.1)	(1.3)	(0.4)	(100.0)				
先天性代謝異常による腎不全	7	13	30	48	29	35	18	11	3	4	198	76	274	165.60	41.50
(%)	(3.5)	(6.6)	(15.2)	(24.2)	(14.6)	(17.7)	(9.1)	(5.6)	(1.5)	(2.0)	(100.0)				
腎・尿路結核	4	14	23	30	31	17	12	3	1	2	137	58	195	159.71	38.48
(%)	(2.9)	(10.2)	(16.8)	(21.9)	(22.6)	(12.4)	(8.8)	(2.2)	(0.7)	(1.5)	(100.0)				
腎・尿路結石	22	41	83	92	88	63	34	8	6	3	440	141	581	156.77	35.80
(%)	(5.0)	(9.3)	(18.9)	(20.9)	(20.0)	(14.3)	(7.7)	(1.8)	(1.4)	(0.7)	(100.0)				
腎・尿路腫瘍	23	76	175	158	110	76	32	15	8	8	681	229	910	152.59	35.26
(%)	(3.4)	(11.2)	(25.7)	(23.2)	(16.2)	(11.2)	(4.7)	(2.2)	(1.2)	(1.2)	(100.0)				
閉塞性尿路障害	9	39	88	119	128	83	40	19	9	5	539	191	730	163.27	35.08
(%)	(1.7)	(7.2)	(16.3)	(22.1)	(23.7)	(15.4)	(7.4)	(3.5)	(1.7)	(0.9)	(100.0)				
骨髄腫	13	17	27	37	42	27	17	8	6	7	201	90	291	164.33	44.10
(%)	(6.5)	(8.5)	(13.4)	(18.4)	(20.9)	(13.4)	(8.5)	(4.0)	(3.0)	(3.5)	(100.0)				
腎形成不全	7	33	98	116	94	80	32	17	12	5	494	166	660	162.94	37.58
(%)	(1.4)	(6.7)	(19.8)	(23.5)	(19.0)	(16.2)	(6.5)	(3.4)	(2.4)	(1.0)	(100.0)				
不明	677	2,120	3,981	4,922	4,191	2,730	1,430	667	308	193	21,219	8,678	29,897	158.06	36.17
(%)	(3.2)	(10.0)	(18.8)	(23.2)	(19.8)	(12.9)	(6.7)	(3.1)	(1.5)	(0.9)	(100.0)				
再導入	40	108	282	360	337	259	128	59	31	22	1,626	557	2,183	163.46	36.48
(%)	(2.5)	(6.6)	(17.3)	(22.1)	(20.7)	(15.9)	(7.9)	(3.6)	(1.9)	(1.4)	(100.0)				
その他	205	571	1,005	1,284	1,102	788	417	223	102	67	5,764	2,280	8,044	159.89	37.99
(%)	(3.6)	(9.9)	(17.4)	(22.3)	(19.1)	(13.7)	(7.2)	(3.9)	(1.8)	(1.2)	(100.0)				
合計	8,409	24,586	45,527	53,667	44,514	28,750	14,958	6,870	2,821	1,860	231,962	81,234	313,196	156.35	35.90
(%)	(3.6)	(10.6)	(19.6)	(23.1)	(19.2)	(12.4)	(6.4)	(3.0)	(1.2)	(0.8)	(100.0)				
記載なし			4		1		2				7	14	21	160.29	38.68
(%)			(57.1)		(14.3)		(28.6)				(100.0)				
総計	8,409	24,586	45,531	53,667	44,515	28,750	14,960	6,870	2,821	1,860	231,969	81,248	313,217	156.35	35.90
(%)	(3.6)	(10.6)	(19.6)	(23.1)	(19.2)	(12.4)	(6.4)	(3.0)	(1.2)	(0.8)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査