

表 3131 総コレステロール濃度(mg/dL) 原疾患別 (施設血液透析患者 透析回数週3回)

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	1,463	4,531	9,269	11,841	10,541	6,974	3,576	1,642	631	393	50,861	12,613	63,474	159.13	35.21
(%)	(2.9)	(8.9)	(18.2)	(23.3)	(20.7)	(13.7)	(7.0)	(3.2)	(1.2)	(0.8)	(100.0)				
慢性腎盂腎炎	45	141	258	371	338	250	131	66	34	13	1,647	393	2,040	162.48	36.36
(%)	(2.7)	(8.6)	(15.7)	(22.5)	(20.5)	(15.2)	(8.0)	(4.0)	(2.1)	(0.8)	(100.0)				
急速進行性糸球体腎炎	26	76	221	274	295	217	148	89	36	42	1,424	347	1,771	171.06	40.96
(%)	(1.8)	(5.3)	(15.5)	(19.2)	(20.7)	(15.2)	(10.4)	(6.3)	(2.5)	(2.9)	(100.0)				
妊娠腎/妊娠中毒症	7	33	91	140	186	161	106	40	18	10	792	189	981	173.60	34.55
(%)	(0.9)	(4.2)	(11.5)	(17.7)	(23.5)	(20.3)	(13.4)	(5.1)	(2.3)	(1.3)	(100.0)				
その他の分類不能の腎炎	24	75	135	169	151	90	60	29	9	9	751	178	929	159.12	37.90
(%)	(3.2)	(10.0)	(18.0)	(22.5)	(20.1)	(12.0)	(8.0)	(3.9)	(1.2)	(1.2)	(100.0)				
多発性嚢胞腎	165	521	1,098	1,567	1,383	919	479	170	75	47	6,424	1,592	8,016	160.01	34.25
(%)	(2.6)	(8.1)	(17.1)	(24.4)	(21.5)	(14.3)	(7.5)	(2.6)	(1.2)	(0.7)	(100.0)				
腎硬化症	555	1,896	3,585	4,359	3,469	2,082	1,039	457	181	99	17,722	4,285	22,007	155.15	34.19
(%)	(3.1)	(10.7)	(20.2)	(24.6)	(19.6)	(11.7)	(5.9)	(2.6)	(1.0)	(0.6)	(100.0)				
悪性高血圧	25	120	283	342	314	192	99	48	21	15	1,459	402	1,861	160.20	35.77
(%)	(1.7)	(8.2)	(19.4)	(23.4)	(21.5)	(13.2)	(6.8)	(3.3)	(1.4)	(1.0)	(100.0)				
糖尿病性腎症	3,702	9,834	16,092	16,573	12,140	7,089	3,430	1,501	588	416	71,365	18,061	89,426	149.73	34.86
(%)	(5.2)	(13.8)	(22.5)	(23.2)	(17.0)	(9.9)	(4.8)	(2.1)	(0.8)	(0.6)	(100.0)				
SLE腎炎	22	88	155	240	247	202	124	60	39	16	1,193	311	1,504	169.23	38.69
(%)	(1.8)	(7.4)	(13.0)	(20.1)	(20.7)	(16.9)	(10.4)	(5.0)	(3.3)	(1.3)	(100.0)				
アミロイド腎	9	25	36	44	43	39	27	7	7	3	240	74	314	163.64	39.93
(%)	(3.8)	(10.4)	(15.0)	(18.3)	(17.9)	(16.3)	(11.3)	(2.9)	(2.9)	(1.3)	(100.0)				
痛風腎	20	72	137	142	122	71	37	21	6	3	631	147	778	155.04	34.25
(%)	(3.2)	(11.4)	(21.7)	(22.5)	(19.3)	(11.3)	(5.9)	(3.3)	(1.0)	(0.5)	(100.0)				
先天性代謝異常による腎不全	7	10	23	32	18	25	7	9	3	3	137	36	173	163.18	44.25
(%)	(5.1)	(7.3)	(16.8)	(23.4)	(13.1)	(18.2)	(5.1)	(6.6)	(2.2)	(2.2)	(100.0)				
腎・尿路結核	4	11	17	24	30	11	8	3		2	110	31	141	159.10	38.76
(%)	(3.6)	(10.0)	(15.5)	(21.8)	(27.3)	(10.0)	(7.3)	(2.7)		(1.8)	(100.0)				
腎・尿路結石	20	29	65	76	64	48	26	4	4	3	339	82	421	155.67	35.64
(%)	(5.9)	(8.6)	(19.2)	(22.4)	(18.9)	(14.2)	(7.7)	(1.2)	(1.2)	(0.9)	(100.0)				
腎・尿路腫瘍	18	67	144	128	82	56	23	10	6	6	540	138	678	150.86	34.50
(%)	(3.3)	(12.4)	(26.7)	(23.7)	(15.2)	(10.4)	(4.3)	(1.9)	(1.1)	(1.1)	(100.0)				
閉塞性尿路障害	6	31	68	87	100	60	24	10	4	4	394	100	494	161.35	33.89
(%)	(1.5)	(7.9)	(17.3)	(22.1)	(25.4)	(15.2)	(6.1)	(2.5)	(1.0)	(1.0)	(100.0)				
骨髄腫	10	11	19	24	32	18	15	7	3	6	145	45	190	166.41	45.44
(%)	(6.9)	(7.6)	(13.1)	(16.6)	(22.1)	(12.4)	(10.3)	(4.8)	(2.1)	(4.1)	(100.0)				
腎形成不全	5	22	73	74	58	47	23	9	5	1	317	83	400	159.02	33.25
(%)	(1.6)	(6.9)	(23.0)	(23.3)	(18.3)	(14.8)	(7.3)	(2.8)	(1.6)	(0.3)	(100.0)				
不明	548	1,705	3,158	3,874	3,165	2,003	1,030	459	209	124	16,275	4,901	21,176	156.45	35.57
(%)	(3.4)	(10.5)	(19.4)	(23.8)	(19.4)	(12.3)	(6.3)	(2.8)	(1.3)	(0.8)	(100.0)				
再導入	29	75	210	238	237	167	80	42	24	10	1,112	244	1,356	162.00	36.04
(%)	(2.6)	(6.7)	(18.9)	(21.4)	(21.3)	(15.0)	(7.2)	(3.8)	(2.2)	(0.9)	(100.0)				
その他	158	465	791	981	835	572	305	162	75	47	4,391	1,324	5,715	158.74	37.78
(%)	(3.6)	(10.6)	(18.0)	(22.3)	(19.0)	(13.0)	(6.9)	(3.7)	(1.7)	(1.1)	(100.0)				
合計	6,868	19,838	35,928	41,600	33,850	21,293	10,797	4,845	1,978	1,272	178,269	45,576	223,845	155.01	35.49
(%)	(3.9)	(11.1)	(20.2)	(23.3)	(19.0)	(11.9)	(6.1)	(2.7)	(1.1)	(0.7)	(100.0)				
記載なし			2								2	5	7	132.50	4.95
(%)			(100.0)								(100.0)				
総計	6,868	19,838	35,930	41,600	33,850	21,293	10,797	4,845	1,978	1,272	178,271	45,581	223,852	155.01	35.49
(%)	(3.9)	(11.1)	(20.2)	(23.3)	(19.0)	(11.9)	(6.1)	(2.7)	(1.1)	(0.7)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査