

表 3142 総コレステロール濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	100未満	100～	120～	140～	160～	180～	200～	220～	240～	260～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	313	1,027	2,399	3,184	2,917	2,113	1,231	599	206	120	14,109	3,353	17,462	163.16	35.74
(%)	(2.2)	(7.3)	(17.0)	(22.6)	(20.7)	(15.0)	(8.7)	(4.2)	(1.5)	(0.9)	(100.0)				
慢性腎盂腎炎	5	29	51	96	103	56	37	19	7	6	409	75	484	166.70	35.43
(%)	(1.2)	(7.1)	(12.5)	(23.5)	(25.2)	(13.7)	(9.0)	(4.6)	(1.7)	(1.5)	(100.0)				
急速進行性糸球体腎炎	4	18	45	61	50	43	34	20	8	6	289	64	353	170.61	39.78
(%)	(1.4)	(6.2)	(15.6)	(21.1)	(17.3)	(14.9)	(11.8)	(6.9)	(2.8)	(2.1)	(100.0)				
妊娠腎/妊娠中毒症	3	7	40	62	74	49	37	28	5	7	312	89	401	175.26	37.18
(%)	(1.0)	(2.2)	(12.8)	(19.9)	(23.7)	(15.7)	(11.9)	(9.0)	(1.6)	(2.2)	(100.0)				
その他の分類不能の腎炎	11	16	32	29	36	33	21	10	5	1	194	67	261	164.20	39.75
(%)	(5.7)	(8.2)	(16.5)	(14.9)	(18.6)	(17.0)	(10.8)	(5.2)	(2.6)	(0.5)	(100.0)				
多発性嚢胞腎	33	99	267	372	339	216	130	55	24	8	1,543	337	1,880	162.20	33.61
(%)	(2.1)	(6.4)	(17.3)	(24.1)	(22.0)	(14.0)	(8.4)	(3.6)	(1.6)	(0.5)	(100.0)				
腎硬化症	80	265	588	793	613	430	203	101	35	19	3,127	693	3,820	158.47	34.40
(%)	(2.6)	(8.5)	(18.8)	(25.4)	(19.6)	(13.8)	(6.5)	(3.2)	(1.1)	(0.6)	(100.0)				
悪性高血圧	4	23	54	67	82	42	28	10	4	6	320	71	391	164.60	35.33
(%)	(1.3)	(7.2)	(16.9)	(20.9)	(25.6)	(13.1)	(8.8)	(3.1)	(1.3)	(1.9)	(100.0)				
糖尿病性腎症	570	1,661	2,939	3,262	2,534	1,470	716	298	121	78	13,649	3,311	16,960	152.26	34.59
(%)	(4.2)	(12.2)	(21.5)	(23.9)	(18.6)	(10.8)	(5.2)	(2.2)	(0.9)	(0.6)	(100.0)				
SLE腎炎	1	18	36	74	70	58	40	17	14	6	334	92	426	173.93	36.55
(%)	(0.3)	(5.4)	(10.8)	(22.2)	(21.0)	(17.4)	(12.0)	(5.1)	(4.2)	(1.8)	(100.0)				
アミロイド腎	1	5	11	7	14	5	4	3	1	1	52	22	74	164.58	43.22
(%)	(1.9)	(9.6)	(21.2)	(13.5)	(26.9)	(9.6)	(7.7)	(5.8)	(1.9)	(1.9)	(100.0)				
痛風腎	7	5	25	35	27	21	9	4	4		137	29	166	160.06	34.75
(%)	(5.1)	(3.6)	(18.2)	(25.5)	(19.7)	(15.3)	(6.6)	(2.9)	(2.9)		(100.0)				
先天性代謝異常による腎不全		2	6	10	7	8	6	2		1	42	7	49	171.12	36.33
(%)		(4.8)	(14.3)	(23.8)	(16.7)	(19.0)	(14.3)	(4.8)		(2.4)	(100.0)				
腎・尿路結核		3	3	4	1	4	2			1	18	13	31	161.39	40.80
(%)		(16.7)	(16.7)	(22.2)	(5.6)	(22.2)	(11.1)			(5.6)	(100.0)				
腎・尿路結石	2	9	15	14	14	8	4	2	1		69	10	79	154.17	35.07
(%)	(2.9)	(13.0)	(21.7)	(20.3)	(20.3)	(11.6)	(5.8)	(2.9)	(1.4)		(100.0)				
腎・尿路腫瘍	3	5	25	26	23	11	5	3	1	1	103	16	119	156.75	34.18
(%)	(2.9)	(4.9)	(24.3)	(25.2)	(22.3)	(10.7)	(4.9)	(2.9)	(1.0)	(1.0)	(100.0)				
閉塞性尿路障害	3		15	20	18	13	10	4	2		85	19	104	166.67	35.56
(%)	(3.5)		(17.6)	(23.5)	(21.2)	(15.3)	(11.8)	(4.7)	(2.4)		(100.0)				
骨髄腫	1	3	3	6	7	3	1		1	1	26	7	33	159.73	38.57
(%)	(3.8)	(11.5)	(11.5)	(23.1)	(26.9)	(11.5)	(3.8)		(3.8)	(3.8)	(100.0)				
腎形成不全	2	8	17	31	27	18	4	2	4	1	114	22	136	161.66	35.00
(%)	(1.8)	(7.0)	(14.9)	(27.2)	(23.7)	(15.8)	(3.5)	(1.8)	(3.5)	(0.9)	(100.0)				
不明	85	312	611	774	721	527	267	124	53	31	3,505	1,198	4,703	161.26	35.62
(%)	(2.4)	(8.9)	(17.4)	(22.1)	(20.6)	(15.0)	(7.6)	(3.5)	(1.5)	(0.9)	(100.0)				
再導入	9	24	59	94	72	75	35	14	5	8	395	88	483	166.19	36.62
(%)	(2.3)	(6.1)	(14.9)	(23.8)	(18.2)	(19.0)	(8.9)	(3.5)	(1.3)	(2.0)	(100.0)				
その他	30	72	160	217	174	148	68	40	15	6	930	230	1,160	161.07	35.56
(%)	(3.2)	(7.7)	(17.2)	(23.3)	(18.7)	(15.9)	(7.3)	(4.3)	(1.6)	(0.6)	(100.0)				
合計	1,167	3,611	7,401	9,238	7,923	5,351	2,892	1,355	517	307	39,762	9,813	49,575	159.08	35.64
(%)	(2.9)	(9.1)	(18.6)	(23.2)	(19.9)	(13.5)	(7.3)	(3.4)	(1.3)	(0.8)	(100.0)				
記載なし			1		1		2				4	2	6	182.50	37.97
(%)			(25.0)		(25.0)		(50.0)				(100.0)				
総計	1,167	3,611	7,402	9,238	7,924	5,351	2,894	1,355	517	307	39,766	9,815	49,581	159.09	35.64
(%)	(2.9)	(9.1)	(18.6)	(23.2)	(19.9)	(13.5)	(7.3)	(3.4)	(1.3)	(0.8)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査