

表 3162 HDL-C濃度(mg/dL) 治療方法(新分類)別 (透析患者全体)

治療方法 (新分類)	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
施設血液透析	735	14,205	41,200	45,715	33,259	19,140	9,447	4,385	1,955	1,816	171,857	76,868	248,725	48.48	16.63
(%)	(0.4)	(8.3)	(24.0)	(26.6)	(19.4)	(11.1)	(5.5)	(2.6)	(1.1)	(1.1)	(100.0)				
Off-line HDF	18	277	746	897	712	430	222	105	51	38	3,496	1,836	5,332	49.75	17.03
(%)	(0.5)	(7.9)	(21.3)	(25.7)	(20.4)	(12.3)	(6.4)	(3.0)	(1.5)	(1.1)	(100.0)				
On-line HDF	87	2,056	6,735	8,229	6,884	4,276	2,345	1,131	541	411	32,695	11,832	44,527	51.06	17.20
(%)	(0.3)	(6.3)	(20.6)	(25.2)	(21.1)	(13.1)	(7.2)	(3.5)	(1.7)	(1.3)	(100.0)				
Push/Pull HDF		3	13	18	20	19	9	3		1	86	24	110	54.29	16.14
(%)		(3.5)	(15.1)	(20.9)	(23.3)	(22.1)	(10.5)	(3.5)		(1.2)	(100.0)				
ハイオフィルトレーション	1	17	43	60	39	25	11	7	2	1	206	61	267	48.72	15.64
(%)	(0.5)	(8.3)	(20.9)	(29.1)	(18.9)	(12.1)	(5.3)	(3.4)	(1.0)	(0.5)	(100.0)				
IHDF	12	149	631	726	536	285	158	78	31	47	2,653	887	3,540	49.79	17.33
(%)	(0.5)	(5.6)	(23.8)	(27.4)	(20.2)	(10.7)	(6.0)	(2.9)	(1.2)	(1.8)	(100.0)				
血液濾過		3	2	2	1						8	8	16	37.00	13.10
(%)		(37.5)	(25.0)	(25.0)	(12.5)						(100.0)				
血液吸着透析 (リクセル等使用)		59	185	282	221	155	84	35	13	11	1,045	331	1,376	51.83	16.69
(%)		(5.6)	(17.7)	(27.0)	(21.1)	(14.8)	(8.0)	(3.3)	(1.2)	(1.1)	(100.0)				
在宅血液透析		14	42	83	81	69	29	14	9	19	360	195	555	57.55	20.22
(%)		(3.9)	(11.7)	(23.1)	(22.5)	(19.2)	(8.1)	(3.9)	(2.5)	(5.3)	(100.0)				
腹膜透析 (CAPD)	12	175	510	619	452	295	155	63	29	29	2,339	2,563	4,902	49.71	17.19
(%)	(0.5)	(7.5)	(21.8)	(26.5)	(19.3)	(12.6)	(6.6)	(2.7)	(1.2)	(1.2)	(100.0)				
腹膜透析 (APD)	4	95	307	331	247	169	82	44	17	15	1,311	1,464	2,775	49.83	17.20
(%)	(0.3)	(7.2)	(23.4)	(25.2)	(18.8)	(12.9)	(6.3)	(3.4)	(1.3)	(1.1)	(100.0)				
腹膜透析 (CCPD)	2	45	163	139	124	77	37	23	8	5	623	469	1,092	49.31	17.11
(%)	(0.3)	(7.2)	(26.2)	(22.3)	(19.9)	(12.4)	(5.9)	(3.7)	(1.3)	(0.8)	(100.0)				
合計	871	17,098	50,577	57,101	42,576	24,940	12,579	5,888	2,656	2,393	216,679	96,538	313,217	48.96	16.78
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.6)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				
記載なし															
(%)															
総計	871	17,098	50,577	57,101	42,576	24,940	12,579	5,888	2,656	2,393	216,679	96,538	313,217	48.96	16.78
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.6)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査