

表 3163 HDL-C濃度(mg/dL) 併用の状況別 (透析患者全体)

併用の状況	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	853	16,771	49,555	55,965	41,716	24,378	12,295	5,752	2,602	2,343	212,230	91,933	304,163	48.94	16.77
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.7)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				
PD単独	12	204	731	846	646	441	229	106	43	41	3,299	3,805	7,104	50.48	17.10
(%)	(0.4)	(6.2)	(22.2)	(25.6)	(19.6)	(13.4)	(6.9)	(3.2)	(1.3)	(1.2)	(100.0)				
HD(F)単独.PDカテあり		5	26	32	27	13	6	3		1	113	69	182	49.04	14.62
(%)		(4.4)	(23.0)	(28.3)	(23.9)	(11.5)	(5.3)	(2.7)		(0.9)	(100.0)				
PD.HD併用週1 HD(F)	6	102	227	213	159	91	42	20	11	7	878	592	1,470	46.83	16.77
(%)	(0.7)	(11.6)	(25.9)	(24.3)	(18.1)	(10.4)	(4.8)	(2.3)	(1.3)	(0.8)	(100.0)				
PD.HD併用週2 HD(F)		9	21	27	14	9	2	4			86	86	172	46.60	14.41
(%)		(10.5)	(24.4)	(31.4)	(16.3)	(10.5)	(2.3)	(4.7)			(100.0)				
PD.HD併用週3 HD(F)		2	7	3	5		1				18	11	29	43.72	12.97
(%)		(11.1)	(38.9)	(16.7)	(27.8)		(5.6)				(100.0)				
PD.HD併用週4 HD(F)															
(%)															
PD.HD併用その他		5	10	15	6	7	2	2		1	48	23	71	50.58	25.53
(%)		(10.4)	(20.8)	(31.3)	(12.5)	(14.6)	(4.2)	(4.2)		(2.1)	(100.0)				
合計	871	17,098	50,577	57,101	42,573	24,939	12,577	5,887	2,656	2,393	216,672	96,519	313,191	48.96	16.78
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.6)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				
不明					3	1	2	1			7	18	25	65.00	12.38
(%)					(42.9)	(14.3)	(28.6)	(14.3)			(100.0)				
記載なし												1	1		
(%)															
総計	871	17,098	50,577	57,101	42,576	24,940	12,579	5,888	2,656	2,393	216,679	96,538	313,217	48.96	16.78
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.6)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査