

表 3167 HDL-C濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	182	3,460	12,669	16,630	14,026	9,032	4,809	2,353	1,048	897	65,106	28,241	93,347	51.76	17.17
(%)	(0.3)	(5.3)	(19.5)	(25.5)	(21.5)	(13.9)	(7.4)	(3.6)	(1.6)	(1.4)	(100.0)				
慢性腎盂腎炎	7	86	355	484	441	310	188	89	35	31	2,026	909	2,935	53.41	17.63
(%)	(0.3)	(4.2)	(17.5)	(23.9)	(21.8)	(15.3)	(9.3)	(4.4)	(1.7)	(1.5)	(100.0)				
急速進行性糸球 体腎炎	8	53	259	374	384	280	161	108	48	41	1,716	762	2,478	56.18	18.94
(%)	(0.5)	(3.1)	(15.1)	(21.8)	(22.4)	(16.3)	(9.4)	(6.3)	(2.8)	(2.4)	(100.0)				
妊娠腎/妊娠中毒 症	2	29	160	241	271	195	125	68	27	16	1,134	411	1,545	56.11	16.85
(%)	(0.2)	(2.6)	(14.1)	(21.3)	(23.9)	(17.2)	(11.0)	(6.0)	(2.4)	(1.4)	(100.0)				
その他の分類不 能の腎炎	1	50	187	253	203	159	74	46	16	9	998	407	1,405	52.20	16.50
(%)	(0.1)	(5.0)	(18.7)	(25.4)	(20.3)	(15.9)	(7.4)	(4.6)	(1.6)	(0.9)	(100.0)				
多発性嚢胞腎	20	428	1,494	2,173	1,879	1,044	564	232	103	88	8,025	3,231	11,256	51.07	16.12
(%)	(0.2)	(5.3)	(18.6)	(27.1)	(23.4)	(13.0)	(7.0)	(2.9)	(1.3)	(1.1)	(100.0)				
腎硬化症	66	1,429	4,711	5,726	4,198	2,408	1,123	514	225	196	20,596	9,209	29,805	48.98	16.17
(%)	(0.3)	(6.9)	(22.9)	(27.8)	(20.4)	(11.7)	(5.5)	(2.5)	(1.1)	(1.0)	(100.0)				
悪性高血圧	6	114	369	519	366	230	102	59	22	24	1,811	807	2,618	50.10	16.92
(%)	(0.3)	(6.3)	(20.4)	(28.7)	(20.2)	(12.7)	(5.6)	(3.3)	(1.2)	(1.3)	(100.0)				
糖尿病性腎症	434	9,420	23,725	22,321	14,163	7,063	3,224	1,414	634	653	83,051	37,227	120,278	45.43	15.69
(%)	(0.5)	(11.3)	(28.6)	(26.9)	(17.1)	(8.5)	(3.9)	(1.7)	(0.8)	(0.8)	(100.0)				
SLE腎炎	3	59	198	322	351	249	165	102	63	31	1,543	684	2,227	57.30	19.07
(%)	(0.2)	(3.8)	(12.8)	(20.9)	(22.7)	(16.1)	(10.7)	(6.6)	(4.1)	(2.0)	(100.0)				
アミロイド腎		16	32	77	69	51	27	15	10	8	305	157	462	56.30	20.37
(%)		(5.2)	(10.5)	(25.2)	(22.6)	(16.7)	(8.9)	(4.9)	(3.3)	(2.6)	(100.0)				
痛風腎	4	54	182	197	156	100	52	20	9	14	788	287	1,075	50.01	18.33
(%)	(0.5)	(6.9)	(23.1)	(25.0)	(19.8)	(12.7)	(6.6)	(2.5)	(1.1)	(1.8)	(100.0)				
先天性代謝異常 による腎不全	2	8	36	36	33	33	17	8	1	5	179	95	274	53.53	18.73
(%)	(1.1)	(4.5)	(20.1)	(20.1)	(18.4)	(18.4)	(9.5)	(4.5)	(0.6)	(2.8)	(100.0)				
腎・尿路結核	1	5	28	34	30	23	10	6	4	3	144	51	195	53.69	18.71
(%)	(0.7)	(3.5)	(19.4)	(23.6)	(20.8)	(16.0)	(6.9)	(4.2)	(2.8)	(2.1)	(100.0)				
腎・尿路結石	1	18	69	129	90	56	27	9	10	3	412	169	581	51.20	16.05
(%)	(0.2)	(4.4)	(16.7)	(31.3)	(21.8)	(13.6)	(6.6)	(2.2)	(2.4)	(0.7)	(100.0)				
腎・尿路腫瘍	5	45	153	188	135	75	32	19	4	5	661	249	910	48.60	16.90
(%)	(0.8)	(6.8)	(23.1)	(28.4)	(20.4)	(11.3)	(4.8)	(2.9)	(0.6)	(0.8)	(100.0)				
閉塞性尿路障害	1	14	98	141	109	79	41	20	4	9	516	214	730	53.02	18.12
(%)	(0.2)	(2.7)	(19.0)	(27.3)	(21.1)	(15.3)	(7.9)	(3.9)	(0.8)	(1.7)	(100.0)				
骨髄腫	2	15	37	44	39	25	14	4	3	3	186	105	291	50.46	17.79
(%)	(1.1)	(8.1)	(19.9)	(23.7)	(21.0)	(13.4)	(7.5)	(2.2)	(1.6)	(1.6)	(100.0)				
腎形成不全	1	21	96	111	105	72	29	18	1	6	460	200	660	51.20	15.86
(%)	(0.2)	(4.6)	(20.9)	(24.1)	(22.8)	(15.7)	(6.3)	(3.9)	(0.2)	(1.3)	(100.0)				
不明	93	1,332	4,315	5,321	4,108	2,494	1,293	551	280	234	20,021	9,876	29,897	49.99	16.85
(%)	(0.5)	(6.7)	(21.6)	(26.6)	(20.5)	(12.5)	(6.5)	(2.8)	(1.4)	(1.2)	(100.0)				
再導入	5	85	290	369	322	211	109	68	25	35	1,519	664	2,183	52.57	18.44
(%)	(0.3)	(5.6)	(19.1)	(24.3)	(21.2)	(13.9)	(7.2)	(4.5)	(1.6)	(2.3)	(100.0)				
その他	27	356	1,112	1,406	1,097	749	393	165	84	82	5,471	2,573	8,044	50.92	17.42
(%)	(0.5)	(6.5)	(20.3)	(25.7)	(20.1)	(13.7)	(7.2)	(3.0)	(1.5)	(1.5)	(100.0)				
合計	871	17,097	50,575	57,096	42,575	24,938	12,579	5,888	2,656	2,393	216,668	96,528	313,196	48.96	16.78
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.6)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				
記載なし		1	2	5	1	2					11	10	21	45.00	11.35
(%)		(9.1)	(18.2)	(45.5)	(9.1)	(18.2)					(100.0)				
総計	871	17,098	50,577	57,101	42,576	24,940	12,579	5,888	2,656	2,393	216,679	96,538	313,217	48.96	16.78
(%)	(0.4)	(7.9)	(23.3)	(26.4)	(19.6)	(11.5)	(5.8)	(2.7)	(1.2)	(1.1)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。

(C)Japanese Society for Dialysis Therapy

2015年調査