

表 3187 HDL-C濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	37	551	2,332	3,249	3,025	2,017	1,186	595	281	206	13,479	3,983	17,462	53.59	17.41
(%)	(0.3)	(4.1)	(17.3)	(24.1)	(22.4)	(15.0)	(8.8)	(4.4)	(2.1)	(1.5)	(100.0)				
慢性腎盂腎炎		15	54	88	82	58	38	21	6	7	369	115	484	55.02	17.66
(%)		(4.1)	(14.6)	(23.8)	(22.2)	(15.7)	(10.3)	(5.7)	(1.6)	(1.9)	(100.0)				
急速進行性糸球体腎炎	2	8	39	61	68	42	37	21	6	5	289	64	353	56.37	18.12
(%)	(0.7)	(2.8)	(13.5)	(21.1)	(23.5)	(14.5)	(12.8)	(7.3)	(2.1)	(1.7)	(100.0)				
妊娠腎/妊娠中毒症	1	3	32	51	75	55	40	22	12	5	296	105	401	59.47	17.36
(%)	(0.3)	(1.0)	(10.8)	(17.2)	(25.3)	(18.6)	(13.5)	(7.4)	(4.1)	(1.7)	(100.0)				
その他の分類不能の腎炎	1	10	37	32	33	41	20	21	2	4	201	60	261	56.12	19.06
(%)	(0.5)	(5.0)	(18.4)	(15.9)	(16.4)	(20.4)	(10.0)	(10.4)	(1.0)	(2.0)	(100.0)				
多発性嚢胞腎	5	74	227	377	389	209	126	40	27	18	1,492	388	1,880	52.62	16.28
(%)	(0.3)	(5.0)	(15.2)	(25.3)	(26.1)	(14.0)	(8.4)	(2.7)	(1.8)	(1.2)	(100.0)				
腎硬化症	6	169	622	812	658	378	178	86	51	35	2,995	825	3,820	50.66	16.84
(%)	(0.2)	(5.6)	(20.8)	(27.1)	(22.0)	(12.6)	(5.9)	(2.9)	(1.7)	(1.2)	(100.0)				
悪性高血圧		20	41	87	62	56	17	8	6	5	302	89	391	52.14	16.57
(%)		(6.6)	(13.6)	(28.8)	(20.5)	(18.5)	(5.6)	(2.6)	(2.0)	(1.7)	(100.0)				
糖尿病性腎症	52	1,330	3,605	3,467	2,341	1,205	548	242	109	91	12,990	3,970	16,960	46.17	15.62
(%)	(0.4)	(10.2)	(27.8)	(26.7)	(18.0)	(9.3)	(4.2)	(1.9)	(0.8)	(0.7)	(100.0)				
SLE腎炎	9	31	53	77	53	40	19	19	20	10	312	114	426	60.50	19.71
(%)		(2.9)	(9.9)	(17.0)	(24.7)	(17.0)	(12.8)	(6.1)	(6.4)	(3.2)	(100.0)				
アミロイド腎	4	8	13	10	6	4	4	7	1	1	54	20	74	55.52	19.71
(%)		(7.4)	(14.8)	(24.1)	(18.5)	(11.1)	(7.4)	(13.0)	(1.9)	(1.9)	(100.0)				
痛風腎	8	30	33	22	20	16	4	4	4	2	139	27	166	52.75	19.30
(%)		(5.8)	(21.6)	(23.7)	(15.8)	(14.4)	(11.5)	(2.9)	(2.9)	(1.4)	(100.0)				
先天性代謝異常による腎不全			2	11	6	9	6	5	1	1	41	8	49	62.51	18.51
(%)			(4.9)	(26.8)	(14.6)	(22.0)	(14.6)	(12.2)	(2.4)	(2.4)	(100.0)				
腎・尿路結核			4	4	9	2	2	2	1	1	25	6	31	59.24	17.82
(%)			(16.0)	(16.0)	(36.0)	(8.0)	(8.0)	(8.0)	(4.0)	(4.0)	(100.0)				
腎・尿路結石		1	5	16	19	8	4	4	4		61	18	79	56.34	16.78
(%)		(1.6)	(8.2)	(26.2)	(31.1)	(13.1)	(6.6)	(6.6)	(6.6)		(100.0)				
腎・尿路腫瘍		5	24	30	17	19	8	1	1		105	14	119	49.43	14.81
(%)		(4.8)	(22.9)	(28.6)	(16.2)	(18.1)	(7.6)	(1.0)	(1.0)		(100.0)				
閉塞性尿路障害		3	14	23	15	16	7	7		1	86	18	104	54.12	16.16
(%)		(3.5)	(16.3)	(26.7)	(17.4)	(18.6)	(8.1)	(8.1)		(1.2)	(100.0)				
骨髄腫			3	7	8	3	1	1	1		24	9	33	54.38	14.22
(%)			(12.5)	(29.2)	(33.3)	(12.5)	(4.2)	(4.2)	(4.2)		(100.0)				
腎形成不全		4	22	27	28	19	7	5		2	114	22	136	52.47	16.01
(%)		(3.5)	(19.3)	(23.7)	(24.6)	(16.7)	(6.1)	(4.4)		(1.8)	(100.0)				
不明	6	168	645	916	770	508	285	129	58	58	3,543	1,160	4,703	52.53	17.56
(%)	(0.2)	(4.7)	(18.2)	(25.9)	(21.7)	(14.3)	(8.0)	(3.6)	(1.6)	(1.6)	(100.0)				
再導入	1	19	59	75	99	59	36	14	6	9	377	106	483	54.10	18.00
(%)	(0.3)	(5.0)	(15.6)	(19.9)	(26.3)	(15.6)	(9.5)	(3.7)	(1.6)	(2.4)	(100.0)				
その他	5	36	155	244	200	127	71	28	10	22	898	262	1,160	52.73	17.67
(%)	(0.6)	(4.0)	(17.3)	(27.2)	(22.3)	(14.1)	(7.9)	(3.1)	(1.1)	(2.4)	(100.0)				
合計	116	2,437	7,991	9,676	8,013	4,910	2,677	1,282	607	483	38,192	11,383	49,575	50.83	17.16
(%)	(0.3)	(6.4)	(20.9)	(25.3)	(21.0)	(12.9)	(7.0)	(3.4)	(1.6)	(1.3)	(100.0)				
記載なし		1	2	3							6		6	38.00	8.25
(%)		(16.7)	(33.3)	(50.0)							(100.0)				
総計	116	2,438	7,993	9,679	8,013	4,910	2,677	1,282	607	483	38,198	11,383	49,581	50.82	17.16
(%)	(0.3)	(6.4)	(20.9)	(25.3)	(21.0)	(12.9)	(7.0)	(3.4)	(1.6)	(1.3)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査