

表 3197 HDL-C濃度(mg/dL) 原疾患別（腹膜透析患者）

原疾患	20未満	20～	30～	40～	50～	60～	70～	80～	90～	100～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	5	82	314	373	284	199	105	47	20	18	1,447	1,383	2,830	50.87	17.31
(%)	(0.3)	(5.7)	(21.7)	(25.8)	(19.6)	(13.8)	(7.3)	(3.2)	(1.4)	(1.2)	(100.0)				
慢性腎盂腎炎		3	8	9	10	15	3		1	1	50	47	97	55.00	21.20
(%)		(6.0)	(16.0)	(18.0)	(20.0)	(30.0)	(6.0)		(2.0)	(2.0)	(100.0)				
急速進行性糸球体腎炎			2	2	6	8	6	5	1		30	36	66	65.43	15.58
(%)			(6.7)	(6.7)	(20.0)	(26.7)	(20.0)	(16.7)	(3.3)		(100.0)				
妊娠腎/妊娠中毒症			5	9	4	1	1				20	7	27	47.60	10.33
(%)			(25.0)	(45.0)	(20.0)	(5.0)	(5.0)				(100.0)				
その他の分類不能の腎炎		2	6	7	12	5	2	3			37	41	78	53.14	15.52
(%)		(5.4)	(16.2)	(18.9)	(32.4)	(13.5)	(5.4)	(8.1)			(100.0)				
多発性嚢胞腎		6	11	30	29	22	11	5	1	1	116	147	263	54.75	15.42
(%)		(5.2)	(9.5)	(25.9)	(25.0)	(19.0)	(9.5)	(4.3)	(0.9)	(0.9)	(100.0)				
腎硬化症	1	26	110	148	107	62	38	20	5	6	523	572	1,095	50.62	16.36
(%)	(0.2)	(5.0)	(21.0)	(28.3)	(20.5)	(11.9)	(7.3)	(3.8)	(1.0)	(1.1)	(100.0)				
悪性高血圧		3	11	17	11	7	3	1	1	2	56	51	107	50.47	17.87
(%)		(5.4)	(19.6)	(30.4)	(19.6)	(12.5)	(5.4)	(1.8)	(1.8)	(3.6)	(100.0)				
糖尿病性腎症	9	152	381	343	220	122	60	29	10	9	1,335	1,464	2,799	45.85	16.09
(%)	(0.7)	(11.4)	(28.5)	(25.7)	(16.5)	(9.1)	(4.5)	(2.2)	(0.7)	(0.7)	(100.0)				
SLE腎炎		2	4	3	5	4	3	3			24	22	46	54.75	18.60
(%)		(8.3)	(16.7)	(12.5)	(20.8)	(16.7)	(12.5)	(12.5)			(100.0)				
アミロイド腎		2		1	1				1		5	10	15	49.40	28.61
(%)		(40.0)		(20.0)	(20.0)				(20.0)		(100.0)				
痛風腎		2	4	1	9	3	1	1		1	22	10	32	55.30	23.32
(%)		(9.1)	(18.2)	(4.5)	(40.9)	(13.6)	(4.5)	(4.5)		(4.5)	(100.0)				
先天性代謝異常による腎不全			1	1	1	3	2			1	9	8	17	63.56	19.76
(%)			(11.1)	(11.1)	(11.1)	(33.3)	(22.2)			(11.1)	(100.0)				
腎・尿路結核								1			1	1	2	87.00	
(%)								(100.0)			(100.0)				
腎・尿路結石			2	5	3	4					14	7	21	50.79	11.20
(%)			(14.3)	(35.7)	(21.4)	(28.6)					(100.0)				
腎・尿路腫瘍			2	4	2	1			1		10	4	14	51.75	17.09
(%)			(20.0)	(40.0)	(20.0)	(10.0)			(10.0)		(100.0)				
閉塞性尿路障害		1	2	1	1	2				1	8	13	21	53.25	27.40
(%)		(12.5)	(25.0)	(12.5)	(12.5)	(25.0)				(12.5)	(100.0)				
骨髄腫		1			1	1					3	4	7	49.00	20.07
(%)		(33.3)			(33.3)	(33.3)					(100.0)				
腎形成不全	1	1	4	5	4	2	1				18	54	72	45.67	15.94
(%)	(5.6)	(5.6)	(22.2)	(27.8)	(22.2)	(11.1)	(5.6)				(100.0)				
不明	1	24	86	103	81	57	32	10	9	5	408	434	842	51.21	17.51
(%)	(0.2)	(5.9)	(21.1)	(25.2)	(19.9)	(14.0)	(7.8)	(2.5)	(2.2)	(1.2)	(100.0)				
再導入		4	4	3	6	3	2	1	2		25	30	55	53.04	21.48
(%)		(16.0)	(16.0)	(12.0)	(24.0)	(12.0)	(8.0)	(4.0)	(8.0)		(100.0)				
その他	1	4	23	24	26	20	4	4	2	4	112	151	263	52.85	18.68
(%)	(0.9)	(3.6)	(20.5)	(21.4)	(23.2)	(17.9)	(3.6)	(3.6)	(1.8)	(3.6)	(100.0)				
合計	18	315	980	1,089	823	541	274	130	54	49	4,273	4,496	8,769	49.69	17.18
(%)	(0.4)	(7.4)	(22.9)	(25.5)	(19.3)	(12.7)	(6.4)	(3.0)	(1.3)	(1.1)	(100.0)				
記載なし															
(%)															
総計	18	315	980	1,089	823	541	274	130	54	49	4,273	4,496	8,769	49.69	17.18
(%)	(0.4)	(7.4)	(22.9)	(25.5)	(19.3)	(12.7)	(6.4)	(3.0)	(1.3)	(1.1)	(100.0)				

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査