

表 3205 Non-HDL-C濃度(mg/dL) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
施設血液透析	2,968	14,263	31,703	37,507	29,991	18,178	9,031	3,861	1,537	989	150,028	98,697	248,725	106.66	33.27
(%)	(2.0)	(9.5)	(21.1)	(25.0)	(20.0)	(12.1)	(6.0)	(2.6)	(1.0)	(0.7)	(100.0)				
Off-line HDF	62	272	611	738	664	353	201	79	34	15	3,029	2,303	5,332	107.47	33.07
(%)	(2.0)	(9.0)	(20.2)	(24.4)	(21.9)	(11.7)	(6.6)	(2.6)	(1.1)	(0.5)	(100.0)				
On-line HDF	434	2,444	5,747	7,194	5,944	3,723	1,924	815	317	150	28,692	15,835	44,527	108.60	32.88
(%)	(1.5)	(8.5)	(20.0)	(25.1)	(20.7)	(13.0)	(6.7)	(2.8)	(1.1)	(0.5)	(100.0)				
Push/Pull HDF	2	9	13	23	13	14	8	1	1	1	85	25	110	109.33	35.42
(%)	(2.4)	(10.6)	(15.3)	(27.1)	(15.3)	(16.5)	(9.4)	(1.2)	(1.2)	(1.2)	(100.0)				
ハイオフィルトレーション	5	11	42	34	42	23	13	7	2	2	181	86	267	110.80	34.79
(%)	(2.8)	(6.1)	(23.2)	(18.8)	(23.2)	(12.7)	(7.2)	(3.9)	(1.1)	(1.1)	(100.0)				
IHDF	46	216	518	545	451	224	131	61	23	9	2,224	1,316	3,540	105.17	32.53
(%)	(2.1)	(9.7)	(23.3)	(24.5)	(20.3)	(10.1)	(5.9)	(2.7)	(1.0)	(0.4)	(100.0)				
血液濾過		1	1	2	1	2					7	9	16	105.00	34.90
(%)		(14.3)	(14.3)	(28.6)	(14.3)	(28.6)					(100.0)				
血液吸着透析 (リクセル等使用)	12	100	175	227	204	132	61	28	11	5	955	421	1,376	108.67	33.03
(%)	(1.3)	(10.5)	(18.3)	(23.8)	(21.4)	(13.8)	(6.4)	(2.9)	(1.2)	(0.5)	(100.0)				
在宅血液透析	2	8	31	72	75	59	40	14	2	5	308	247	555	121.89	32.39
(%)	(0.6)	(2.6)	(10.1)	(23.4)	(24.4)	(19.2)	(13.0)	(4.5)	(0.6)	(1.6)	(100.0)				
腹膜透析 (CAPD)	11	57	178	353	442	336	269	154	69	55	1,924	2,978	4,902	130.22	37.78
(%)	(0.6)	(3.0)	(9.3)	(18.3)	(23.0)	(17.5)	(14.0)	(8.0)	(3.6)	(2.9)	(100.0)				
腹膜透析 (APD)	6	45	127	221	272	183	131	79	35	23	1,122	1,653	2,775	125.20	37.57
(%)	(0.5)	(4.0)	(11.3)	(19.7)	(24.2)	(16.3)	(11.7)	(7.0)	(3.1)	(2.0)	(100.0)				
腹膜透析 (CCPD)	1	26	62	116	113	99	63	25	15	12	532	560	1,092	124.43	37.60
(%)	(0.2)	(4.9)	(11.7)	(21.8)	(21.2)	(18.6)	(11.8)	(4.7)	(2.8)	(2.3)	(100.0)				
合計	3,549	17,452	39,208	47,032	38,212	23,326	11,872	5,124	2,046	1,266	189,087	124,130	313,217	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				
記載なし															
(%)															
総計	3,549	17,452	39,208	47,032	38,212	23,326	11,872	5,124	2,046	1,266	189,087	124,130	313,217	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査