

表 3206 Non-HDL-C濃度(mg/dL) 併用の状況別（透析患者全体）

併用の状況	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	3,530	17,311	38,816	46,304	37,356	22,688	11,397	4,861	1,925	1,176	185,364	118,799	304,163	106.99	33.21
(%)	(1.9)	(9.3)	(20.9)	(25.0)	(20.2)	(12.2)	(6.1)	(2.6)	(1.0)	(0.6)	(100.0)				
PD単独	10	81	261	500	616	496	363	213	102	80	2,722	4,382	7,104	130.17	38.30
(%)	(0.4)	(3.0)	(9.6)	(18.4)	(22.6)	(18.2)	(13.3)	(7.8)	(3.7)	(2.9)	(100.0)				
HD(F)単独.PDカテあり	1	10	18	25	19	11	5	1	1		91	91	182	104.52	30.45
(%)	(1.1)	(11.0)	(19.8)	(27.5)	(20.9)	(12.1)	(5.5)	(1.1)	(1.1)		(100.0)				
PD.HD併用週1 HD(F)	6	44	95	170	188	111	89	38	15	8	764	706	1,470	119.89	34.77
(%)	(0.8)	(5.8)	(12.4)	(22.3)	(24.6)	(14.5)	(11.6)	(5.0)	(2.0)	(1.0)	(100.0)				
PD.HD併用週2 HD(F)		3	7	17	21	12	10	7	3	1	81	91	172	127.05	35.05
(%)		(3.7)	(8.6)	(21.0)	(25.9)	(14.8)	(12.3)	(8.6)	(3.7)	(1.2)	(100.0)				
PD.HD併用週3 HD(F)		1	2	4	4	2	2	2			17	12	29	119.47	36.02
(%)		(5.9)	(11.8)	(23.5)	(23.5)	(11.8)	(11.8)	(11.8)			(100.0)				
PD.HD併用週4 HD(F)															
(%)															
PD.HD併用その他	2	2	7	11	8	5	4	1		1	41	30	71	111.10	35.35
(%)	(4.9)	(4.9)	(17.1)	(26.8)	(19.5)	(12.2)	(9.8)	(2.4)		(2.4)	(100.0)				
合計	3,549	17,452	39,206	47,031	38,212	23,325	11,870	5,123	2,046	1,266	189,080	124,111	313,191	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				
不明			2	1		1	2	1			7	18	25	126.00	40.64
(%)			(28.6)	(14.3)		(14.3)	(28.6)	(14.3)			(100.0)				
記載なし												1	1		
(%)															
総計	3,549	17,452	39,208	47,032	38,212	23,326	11,872	5,124	2,046	1,266	189,087	124,130	313,217	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査