

表 3208 Non-HDL-C濃度(mg/dL) 年齢(歳)別 (透析患者全体)

年齢	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
15歳未満	1		4	3	2	1	1	1	1	2	16	87	103	129.19	60.11
(%)	(6.3)		(25.0)	(18.8)	(12.5)	(6.3)	(6.3)	(6.3)	(6.3)	(12.5)	(100.0)				
15歳～	11	57	129	134	105	57	23	9	3	5	533	437	970	102.91	33.07
(%)	(2.1)	(10.7)	(24.2)	(25.1)	(19.7)	(10.7)	(4.3)	(1.7)	(0.6)	(0.9)	(100.0)				
30歳～	104	662	1,550	2,066	1,811	1,092	620	292	112	89	8,398	5,458	13,856	111.44	34.58
(%)	(1.2)	(7.9)	(18.5)	(24.6)	(21.6)	(13.0)	(7.4)	(3.5)	(1.3)	(1.1)	(100.0)				
45歳～	591	2,690	6,222	8,085	7,169	4,864	2,694	1,212	499	319	34,345	21,996	56,341	111.77	34.89
(%)	(1.7)	(7.8)	(18.1)	(23.5)	(20.9)	(14.2)	(7.8)	(3.5)	(1.5)	(0.9)	(100.0)				
60歳～	1,648	7,954	17,706	21,229	17,392	10,827	5,467	2,408	939	558	86,128	55,506	141,634	107.58	33.51
(%)	(1.9)	(9.2)	(20.6)	(24.6)	(20.2)	(12.6)	(6.3)	(2.8)	(1.1)	(0.6)	(100.0)				
75歳～	1,141	5,747	12,863	14,623	11,156	6,139	2,944	1,141	468	282	56,504	38,244	94,748	104.14	31.91
(%)	(2.0)	(10.2)	(22.8)	(25.9)	(19.7)	(10.9)	(5.2)	(2.0)	(0.8)	(0.5)	(100.0)				
90歳～	53	342	734	891	577	346	123	61	24	11	3,162	2,398	5,560	102.55	30.55
(%)	(1.7)	(10.8)	(23.2)	(28.2)	(18.2)	(10.9)	(3.9)	(1.9)	(0.8)	(0.3)	(100.0)				
合計	3,549	17,452	39,208	47,031	38,212	23,326	11,872	5,124	2,046	1,266	189,086	124,126	313,212	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				
不明				1							1	3	4	100.00	
(%)				(100.0)							(100.0)				
記載なし												1	1		
(%)															
総計	3,549	17,452	39,208	47,032	38,212	23,326	11,872	5,124	2,046	1,266	189,087	124,130	313,217	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				
平均	68.71	68.88	68.72	68.19	67.44	66.71	65.87	65.30	65.28	64.59	67.76	68.01	67.86		
標準偏差	11.91	12.18	12.31	12.45	12.42	12.31	12.32	12.33	12.53	12.99	12.40	12.63	12.49		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査