

表 3210 Non-HDL-C濃度(mg/dL) 原疾患別（透析患者全体）

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	917	4,767	11,265	14,078	12,112	7,548	3,775	1,611	618	331	57,022	36,325	93,347	108.84	32.95
(%)	(1.6)	(8.4)	(19.8)	(24.7)	(21.2)	(13.2)	(6.6)	(2.8)	(1.1)	(0.6)	(100.0)				
慢性腎盂腎炎	31	153	307	421	373	259	116	60	22	14	1,756	1,179	2,935	110.24	33.96
(%)	(1.8)	(8.7)	(17.5)	(24.0)	(21.2)	(14.7)	(6.6)	(3.4)	(1.3)	(0.8)	(100.0)				
急速進行性糸球体腎炎	20	90	249	361	308	211	117	71	26	28	1,481	997	2,478	115.47	36.97
(%)	(1.4)	(6.1)	(16.8)	(24.4)	(20.8)	(14.2)	(7.9)	(4.8)	(1.8)	(1.9)	(100.0)				
妊娠腎/妊娠中毒症	9	48	139	221	249	163	85	45	16	10	985	560	1,545	118.06	33.34
(%)	(0.9)	(4.9)	(14.1)	(22.4)	(25.3)	(16.5)	(8.6)	(4.6)	(1.6)	(1.0)	(100.0)				
その他の分類不能の腎炎	14	66	167	224	152	117	50	33	11	6	840	565	1,405	109.56	33.75
(%)	(1.7)	(7.9)	(19.9)	(26.7)	(18.1)	(13.9)	(6.0)	(3.9)	(1.3)	(0.7)	(100.0)				
多発性嚢胞腎	103	499	1,348	1,741	1,645	897	455	199	76	38	7,001	4,255	11,256	109.78	32.20
(%)	(1.5)	(7.1)	(19.3)	(24.9)	(23.5)	(12.8)	(6.5)	(2.8)	(1.1)	(0.5)	(100.0)				
腎硬化症	306	1,536	3,751	4,692	3,717	2,184	1,112	475	179	95	18,047	11,758	29,805	107.40	32.42
(%)	(1.7)	(8.5)	(20.8)	(26.0)	(20.6)	(12.1)	(6.2)	(2.6)	(1.0)	(0.5)	(100.0)				
悪性高血圧	24	104	308	389	320	215	116	53	25	15	1,569	1,049	2,618	111.45	34.41
(%)	(1.5)	(6.6)	(19.6)	(24.8)	(20.4)	(13.7)	(7.4)	(3.4)	(1.6)	(1.0)	(100.0)				
糖尿病性腎症	1,639	7,836	16,163	18,045	13,661	8,169	4,198	1,787	752	485	72,735	47,543	120,278	104.98	33.64
(%)	(2.3)	(10.8)	(22.2)	(24.8)	(18.8)	(11.2)	(5.8)	(2.5)	(1.0)	(0.7)	(100.0)				
SLE腎炎	16	104	194	353	305	208	119	47	25	11	1,382	845	2,227	114.35	34.42
(%)	(1.2)	(7.5)	(14.0)	(25.5)	(22.1)	(15.1)	(8.6)	(3.4)	(1.8)	(0.8)	(100.0)				
アミロイド腎	13	17	54	55	52	31	16	12	1	3	254	208	462	107.78	37.99
(%)	(5.1)	(6.7)	(21.3)	(21.7)	(20.5)	(12.2)	(6.3)	(4.7)	(0.4)	(1.2)	(100.0)				
痛風腎	16	64	142	166	140	74	57	20	8	2	689	386	1,075	107.70	33.35
(%)	(2.3)	(9.3)	(20.6)	(24.1)	(20.3)	(10.7)	(8.3)	(2.9)	(1.2)	(0.3)	(100.0)				
先天性代謝異常による腎不全	4	12	32	31	40	18	9	6	2	2	156	118	274	111.04	38.07
(%)	(2.6)	(7.7)	(20.5)	(19.9)	(25.6)	(11.5)	(5.8)	(3.8)	(1.3)	(1.3)	(100.0)				
腎・尿路結核	4	15	24	21	30	17	8	1		1	121	74	195	104.64	33.77
(%)	(3.3)	(12.4)	(19.8)	(17.4)	(24.8)	(14.0)	(6.6)	(0.8)		(0.8)	(100.0)				
腎・尿路結石	10	33	84	74	71	49	23	8	2	1	355	226	581	105.38	32.93
(%)	(2.8)	(9.3)	(23.7)	(20.8)	(20.0)	(13.8)	(6.5)	(2.3)	(0.6)	(0.3)	(100.0)				
腎・尿路腫瘍	10	56	134	162	100	65	25	15	6	7	580	330	910	104.92	33.84
(%)	(1.7)	(9.7)	(23.1)	(27.9)	(17.2)	(11.2)	(4.3)	(2.6)	(1.0)	(1.2)	(100.0)				
閉塞性尿路障害	5	36	81	115	110	53	31	16	2	4	453	277	730	110.32	31.91
(%)	(1.1)	(7.9)	(17.9)	(25.4)	(24.3)	(11.7)	(6.8)	(3.5)	(0.4)	(0.9)	(100.0)				
骨髄腫	3	19	26	30	34	20	17	6	3	5	163	128	291	114.61	41.61
(%)	(1.8)	(11.7)	(16.0)	(18.4)	(20.9)	(12.3)	(10.4)	(3.7)	(1.8)	(3.1)	(100.0)				
腎形成不全	5	29	75	91	105	50	25	8	7	3	398	262	660	110.14	32.82
(%)	(1.3)	(7.3)	(18.8)	(22.9)	(26.4)	(12.6)	(6.3)	(2.0)	(1.8)	(0.8)	(100.0)				
不明	287	1,462	3,515	4,253	3,470	2,199	1,084	472	184	150	17,076	12,821	29,897	108.24	33.59
(%)	(1.7)	(8.6)	(20.6)	(24.9)	(20.3)	(12.9)	(6.3)	(2.8)	(1.1)	(0.9)	(100.0)				
再導入	21	107	227	331	273	194	106	40	22	15	1,336	847	2,183	111.86	34.83
(%)	(1.6)	(8.0)	(17.0)	(24.8)	(20.4)	(14.5)	(7.9)	(3.0)	(1.6)	(1.1)	(100.0)				
その他	92	399	920	1,178	945	584	327	139	58	40	4,682	3,362	8,044	108.95	34.58
(%)	(2.0)	(8.5)	(19.6)	(25.2)	(20.2)	(12.5)	(7.0)	(3.0)	(1.2)	(0.9)	(100.0)				
合計	3,549	17,452	39,205	47,032	38,212	23,325	11,871	5,124	2,045	1,266	189,081	124,115	313,196	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				
記載なし			3			1	1		1		6	15	21	122.17	47.60
(%)			(50.0)			(16.7)	(16.7)		(16.7)		(100.0)				
総計	3,549	17,452	39,208	47,032	38,212	23,326	11,872	5,124	2,046	1,266	189,087	124,130	313,217	107.39	33.42
(%)	(1.9)	(9.2)	(20.7)	(24.9)	(20.2)	(12.3)	(6.3)	(2.7)	(1.1)	(0.7)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査