

表 3219 Non-HDL-C濃度(mg/dL) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	707	3,591	8,316	10,306	8,726	5,302	2,605	1,105	418	226	41,302	22,172	63,474	107.97	32.74
(%)	(1.7)	(8.7)	(20.1)	(25.0)	(21.1)	(12.8)	(6.3)	(2.7)	(1.0)	(0.5)	(100.0)				
慢性腎盂腎炎	27	119	232	306	265	196	78	39	16	9	1,287	753	2,040	108.92	33.87
(%)	(2.1)	(9.2)	(18.0)	(23.8)	(20.6)	(15.2)	(6.1)	(3.0)	(1.2)	(0.7)	(100.0)				
急速進行性糸球体腎炎	18	68	201	273	240	166	90	51	14	21	1,142	629	1,771	114.54	36.71
(%)	(1.6)	(6.0)	(17.6)	(23.9)	(21.0)	(14.5)	(7.9)	(4.5)	(1.2)	(1.8)	(100.0)				
妊娠腎/妊娠中毒症	6	33	97	140	161	114	59	26	11	7	654	327	981	117.98	33.38
(%)	(0.9)	(5.0)	(14.8)	(21.4)	(24.6)	(17.4)	(9.0)	(4.0)	(1.7)	(1.1)	(100.0)				
その他の分類不能の腎炎	9	49	129	168	102	88	32	23	9	4	613	316	929	108.86	33.54
(%)	(1.5)	(8.0)	(21.0)	(27.4)	(16.6)	(14.4)	(5.2)	(3.8)	(1.5)	(0.7)	(100.0)				
多発性嚢胞腎	86	387	1,002	1,329	1,241	681	340	145	49	28	5,288	2,728	8,016	109.33	32.04
(%)	(1.6)	(7.3)	(18.9)	(25.1)	(23.5)	(12.9)	(6.4)	(2.7)	(0.9)	(0.5)	(100.0)				
腎硬化症	245	1,255	3,049	3,731	2,900	1,685	827	348	136	67	14,243	7,764	22,007	106.50	32.01
(%)	(1.7)	(8.8)	(21.4)	(26.2)	(20.4)	(11.8)	(5.8)	(2.4)	(1.0)	(0.5)	(100.0)				
悪性高血圧	17	86	236	297	247	160	84	40	18	11	1,196	665	1,861	110.79	34.22
(%)	(1.4)	(7.2)	(19.7)	(24.8)	(20.7)	(13.4)	(7.0)	(3.3)	(1.5)	(0.9)	(100.0)				
糖尿病性腎症	1,335	6,407	13,048	14,326	10,726	6,433	3,222	1,353	580	378	57,808	31,618	89,426	104.38	33.49
(%)	(2.3)	(11.1)	(22.6)	(24.8)	(18.6)	(11.1)	(5.6)	(2.3)	(1.0)	(0.7)	(100.0)				
SLE腎炎	14	83	141	261	207	153	86	33	17	6	1,001	503	1,504	113.15	34.37
(%)	(1.4)	(8.3)	(14.1)	(26.1)	(20.7)	(15.3)	(8.6)	(3.3)	(1.7)	(0.6)	(100.0)				
アミロイド腎	11	15	44	37	43	22	10	10	1	2	195	119	314	106.27	36.91
(%)	(5.6)	(7.7)	(22.6)	(19.0)	(22.1)	(11.3)	(5.1)	(5.1)	(0.5)	(1.0)	(100.0)				
痛風腎	11	49	110	124	109	54	39	14	7		517	261	778	106.85	32.62
(%)	(2.1)	(9.5)	(21.3)	(24.0)	(21.1)	(10.4)	(7.5)	(2.7)	(1.4)		(100.0)				
先天性代謝異常による腎不全	4	11	22	23	19	12	6	5	1	2	105	68	173	109.31	42.04
(%)	(3.8)	(10.5)	(21.0)	(21.9)	(18.1)	(11.4)	(5.7)	(4.8)	(1.0)	(1.9)	(100.0)				
腎・尿路結核	3	12	19	17	22	14	7	1		1	96	45	141	105.30	35.16
(%)	(3.1)	(12.5)	(19.8)	(17.7)	(22.9)	(14.6)	(7.3)	(1.0)		(1.0)	(100.0)				
腎・尿路結石	8	24	61	65	50	37	20	4	1	1	271	150	421	105.30	32.78
(%)	(3.0)	(8.9)	(22.5)	(24.0)	(18.5)	(13.7)	(7.4)	(1.5)	(0.4)	(0.4)	(100.0)				
腎・尿路腫瘍	7	50	108	126	77	51	16	12	5	4	456	222	678	103.74	33.03
(%)	(1.5)	(11.0)	(23.7)	(27.6)	(16.9)	(11.2)	(3.5)	(2.6)	(1.1)	(0.9)	(100.0)				
閉塞性尿路障害	5	27	58	89	78	40	23	7		2	329	165	494	108.24	30.37
(%)	(1.5)	(8.2)	(17.6)	(27.1)	(23.7)	(12.2)	(7.0)	(2.1)		(0.6)	(100.0)				
骨髄腫	1	13	17	22	26	14	15	4	2	5	119	71	190	117.87	42.13
(%)	(0.8)	(10.9)	(14.3)	(18.5)	(21.8)	(11.8)	(12.6)	(3.4)	(1.7)	(4.2)	(100.0)				
腎形成不全	3	21	50	61	69	29	16	7	3	1	260	140	400	108.28	31.40
(%)	(1.2)	(8.1)	(19.2)	(23.5)	(26.5)	(11.2)	(6.2)	(2.7)	(1.2)	(0.4)	(100.0)				
不明	227	1,116	2,720	3,304	2,651	1,627	772	324	121	107	12,969	8,207	21,176	107.29	32.94
(%)	(1.8)	(8.6)	(21.0)	(25.5)	(20.4)	(12.5)	(6.0)	(2.5)	(0.9)	(0.8)	(100.0)				
再導入	17	73	168	219	172	141	58	27	14	11	900	456	1,356	110.47	35.22
(%)	(1.9)	(8.1)	(18.7)	(24.3)	(19.1)	(15.7)	(6.4)	(3.0)	(1.6)	(1.2)	(100.0)				
その他	73	302	713	886	716	425	245	111	42	27	3,540	2,175	5,715	108.58	34.51
(%)	(2.1)	(8.5)	(20.1)	(25.0)	(20.2)	(12.0)	(6.9)	(3.1)	(1.2)	(0.8)	(100.0)				
合計	2,834	13,791	30,541	36,110	28,847	17,444	8,650	3,689	1,465	920	144,291	79,554	223,845	106.56	33.18
(%)	(2.0)	(9.6)	(21.2)	(25.0)	(20.0)	(12.1)	(6.0)	(2.6)	(1.0)	(0.6)	(100.0)				
記載なし			1								1	6	7	76.00	
(%)			(100.0)								(100.0)				
総計	2,834	13,791	30,542	36,110	28,847	17,444	8,650	3,689	1,465	920	144,292	79,560	223,852	106.56	33.18
(%)	(2.0)	(9.6)	(21.2)	(25.0)	(20.0)	(12.1)	(6.0)	(2.6)	(1.0)	(0.6)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査