

表 3228 Non-HDL-C濃度(mg/dL) 原疾患別（血液透析濾過患者 透析回数週3回）

原疾患	50未満	50～	70～	90～	110～	130～	150～	170～	190～	210～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	168	927	2,335	2,920	2,547	1,657	819	329	129	55	11,886	5,576	17,462	109.50	32.51
(%)	(1.4)	(7.8)	(19.6)	(24.6)	(21.4)	(13.9)	(6.9)	(2.8)	(1.1)	(0.5)	(100.0)				
慢性腎盂腎炎	4	26	54	82	83	46	25	12	5	1	338	146	484	111.99	32.23
(%)	(1.2)	(7.7)	(16.0)	(24.3)	(24.6)	(13.6)	(7.4)	(3.6)	(1.5)	(0.3)	(100.0)				
急速進行性糸球体腎炎	2	17	42	72	51	25	17	14	5	3	248	105	353	112.97	35.51
(%)	(0.8)	(6.9)	(16.9)	(29.0)	(20.6)	(10.1)	(6.9)	(5.6)	(2.0)	(1.2)	(100.0)				
妊娠腎/妊娠中毒症	3	11	40	60	73	35	21	14	4	1	262	139	401	116.39	32.08
(%)	(1.1)	(4.2)	(15.3)	(22.9)	(27.9)	(13.4)	(8.0)	(5.3)	(1.5)	(0.4)	(100.0)				
その他の分類不能の腎炎	4	14	29	38	36	19	6	6	2	1	155	106	261	107.56	33.71
(%)	(2.6)	(9.0)	(18.7)	(24.5)	(23.2)	(12.3)	(3.9)	(3.9)	(1.3)	(0.6)	(100.0)				
多発性嚢胞腎	12	95	280	317	315	164	81	34	18	2	1,318	562	1,880	109.21	30.60
(%)	(0.9)	(7.2)	(21.2)	(24.1)	(23.9)	(12.4)	(6.1)	(2.6)	(1.4)	(0.2)	(100.0)				
腎硬化症	42	213	520	705	555	326	169	72	23	13	2,638	1,182	3,820	108.08	32.16
(%)	(1.6)	(8.1)	(19.7)	(26.7)	(21.0)	(12.4)	(6.4)	(2.7)	(0.9)	(0.5)	(100.0)				
悪性高血圧	6	13	57	61	50	41	20	11	4	1	264	127	391	111.48	33.63
(%)	(2.3)	(4.9)	(21.6)	(23.1)	(18.9)	(15.5)	(7.6)	(4.2)	(1.5)	(0.4)	(100.0)				
糖尿病性腎症	233	1,126	2,448	2,886	2,235	1,279	696	287	116	56	11,362	5,598	16,960	105.81	32.97
(%)	(2.1)	(9.9)	(21.5)	(25.4)	(19.7)	(11.3)	(6.1)	(2.5)	(1.0)	(0.5)	(100.0)				
SLE腎炎	1	20	42	71	68	39	21	8	5	2	277	149	426	113.95	32.62
(%)	(0.4)	(7.2)	(15.2)	(25.6)	(24.5)	(14.1)	(7.6)	(2.9)	(1.8)	(0.7)	(100.0)				
アミロイド腎	1	2	6	16	5	7	4	1		1	43	31	74	113.35	42.95
(%)	(2.3)	(4.7)	(14.0)	(37.2)	(11.6)	(16.3)	(9.3)	(2.3)		(2.3)	(100.0)				
痛風腎	2	13	27	31	22	13	10	6	1	1	126	40	166	108.17	35.26
(%)	(1.6)	(10.3)	(21.4)	(24.6)	(17.5)	(10.3)	(7.9)	(4.8)	(0.8)	(0.8)	(100.0)				
先天性代謝異常による腎不全		1	10	4	13	4	2	1	1		36	13	49	111.69	32.02
(%)		(2.8)	(27.8)	(11.1)	(36.1)	(11.1)	(5.6)	(2.8)	(2.8)		(100.0)				
腎・尿路結核	1	3	4	2	4	2	1				17	14	31	97.29	31.39
(%)	(5.9)	(17.6)	(23.5)	(11.8)	(23.5)	(11.8)	(5.9)				(100.0)				
腎・尿路結石	2	8	17	8	11	6	1	2			55	24	79	97.64	31.10
(%)	(3.6)	(14.5)	(30.9)	(14.5)	(20.0)	(10.9)	(1.8)	(3.6)			(100.0)				
腎・尿路腫瘍	2	3	23	27	20	9	5	2		2	93	26	119	107.03	33.26
(%)	(2.2)	(3.2)	(24.7)	(29.0)	(21.5)	(9.7)	(5.4)	(2.2)		(2.2)	(100.0)				
閉塞性尿路障害		4	14	17	19	10	5	6			75	29	104	115.21	30.40
(%)		(5.3)	(18.7)	(22.7)	(25.3)	(13.3)	(6.7)	(8.0)			(100.0)				
骨髄腫	1	4	4	5	2	3	1	2			22	11	33	103.27	40.92
(%)	(4.5)	(18.2)	(18.2)	(22.7)	(9.1)	(13.6)	(4.5)	(9.1)			(100.0)				
腎形成不全	2	8	19	23	26	12	6	1	3	1	101	35	136	109.97	34.24
(%)	(2.0)	(7.9)	(18.8)	(22.8)	(25.7)	(11.9)	(5.9)	(1.0)	(3.0)	(1.0)	(100.0)				
不明	37	277	604	688	584	377	209	93	28	19	2,916	1,787	4,703	108.65	33.62
(%)	(1.3)	(9.5)	(20.7)	(23.6)	(20.0)	(12.9)	(7.2)	(3.2)	(1.0)	(0.7)	(100.0)				
再導入	3	27	52	84	71	42	38	9	6	3	335	148	483	113.76	34.04
(%)	(0.9)	(8.1)	(15.5)	(25.1)	(21.2)	(12.5)	(11.3)	(2.7)	(1.8)	(0.9)	(100.0)				
その他	12	74	146	213	146	100	58	17	10	5	781	379	1,160	107.96	32.86
(%)	(1.5)	(9.5)	(18.7)	(27.3)	(18.7)	(12.8)	(7.4)	(2.2)	(1.3)	(0.6)	(100.0)				
合計	538	2,886	6,773	8,330	6,936	4,216	2,215	927	360	167	33,348	16,227	49,575	108.18	32.81
(%)	(1.6)	(8.7)	(20.3)	(25.0)	(20.8)	(12.6)	(6.6)	(2.8)	(1.1)	(0.5)	(100.0)				
記載なし			1			1	1		1		4	2	6	144.25	42.69
(%)			(25.0)			(25.0)	(25.0)		(25.0)		(100.0)				
総計	538	2,886	6,774	8,330	6,936	4,217	2,216	927	361	167	33,352	16,229	49,581	108.18	32.81
(%)	(1.6)	(8.7)	(20.3)	(25.0)	(20.8)	(12.6)	(6.6)	(2.8)	(1.1)	(0.5)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査