

表 3244 透析前収縮期血圧(mmHg) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
施設血液透析	33	327	2,794	15,448	46,841	71,941	53,752	20,903	5,280	1,093	218,412	30,313	248,725	151.80	24.28
(%)	(0.0)	(0.1)	(1.3)	(7.1)	(21.4)	(32.9)	(24.6)	(9.6)	(2.4)	(0.5)	(100.0)				
Off-line HDF	3	13	163	508	1,032	1,433	1,012	417	88	18	4,687	645	5,332	147.26	26.32
(%)	(0.1)	(0.3)	(3.5)	(10.8)	(22.0)	(30.6)	(21.6)	(8.9)	(1.9)	(0.4)	(100.0)				
On-line HDF	7	69	623	2,783	8,337	12,866	9,752	3,918	1,061	247	39,663	4,864	44,527	152.20	24.91
(%)	(0.0)	(0.2)	(1.6)	(7.0)	(21.0)	(32.4)	(24.6)	(9.9)	(2.7)	(0.6)	(100.0)				
Push/Pull HDF			2	8	17	35	24	11	3	1	101	9	110	153.06	25.45
(%)			(2.0)	(7.9)	(16.8)	(34.7)	(23.8)	(10.9)	(3.0)	(1.0)	(100.0)				
ハイオフィルトレーション	1	4	17	45	50	60	39	16	11	1	244	23	267	141.53	32.86
(%)	(0.4)	(1.6)	(7.0)	(18.4)	(20.5)	(24.6)	(16.0)	(6.6)	(4.5)	(0.4)	(100.0)				
IHDF		8	55	266	705	1,040	811	347	93	14	3,339	201	3,540	151.84	25.31
(%)		(0.2)	(1.6)	(8.0)	(21.1)	(31.1)	(24.3)	(10.4)	(2.8)	(0.4)	(100.0)				
血液濾過			1	2	3	3	1	1			11	5	16	137.64	28.10
(%)			(9.1)	(18.2)	(27.3)	(27.3)	(9.1)	(9.1)			(100.0)				
血液吸着透析 (リクセル等使用)	2	4	41	163	329	413	223	70	10	3	1,258	118	1,376	142.76	24.69
(%)	(0.2)	(0.3)	(3.3)	(13.0)	(26.2)	(32.8)	(17.7)	(5.6)	(0.8)	(0.2)	(100.0)				
在宅血液透析			13	54	105	126	46	10	3	1	358	197	555	139.47	22.51
(%)			(3.6)	(15.1)	(29.3)	(35.2)	(12.8)	(2.8)	(0.8)	(0.3)	(100.0)				
腹膜透析 (CAPD)		9	99	343	969	924	434	113	23	5	2,919	1,983	4,902	140.74	22.74
(%)		(0.3)	(3.4)	(11.8)	(33.2)	(31.7)	(14.9)	(3.9)	(0.8)	(0.2)	(100.0)				
腹膜透析 (APD)		8	55	208	544	553	232	54	12	3	1,669	1,106	2,775	139.95	22.35
(%)		(0.5)	(3.3)	(12.5)	(32.6)	(33.1)	(13.9)	(3.2)	(0.7)	(0.2)	(100.0)				
腹膜透析 (CCPD)		3	15	69	210	237	134	42	12	3	725	367	1,092	145.39	23.78
(%)		(0.4)	(2.1)	(9.5)	(29.0)	(32.7)	(18.5)	(5.8)	(1.7)	(0.4)	(100.0)				
合計	46	445	3,878	19,897	59,142	89,631	66,460	25,902	6,596	1,389	273,386	39,831	313,217	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				
記載なし															
(%)															
総計	46	445	3,878	19,897	59,142	89,631	66,460	25,902	6,596	1,389	273,386	39,831	313,217	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査