

表 3245 透析前収縮期血圧(mmHg) 併用の状況別（透析患者全体）

併用の状況	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
HD(F)単独	46	424	3,704	19,262	57,354	87,848	65,603	25,673	6,542	1,375	267,831	36,332	304,163	151.71	24.45
(%)	(0.0)	(0.2)	(1.4)	(7.2)	(21.4)	(32.8)	(24.5)	(9.6)	(2.4)	(0.5)	(100.0)				
PD単独		16	141	512	1,379	1,267	517	113	23	4	3,972	3,132	7,104	138.82	21.95
(%)		(0.4)	(3.5)	(12.9)	(34.7)	(31.9)	(13.0)	(2.8)	(0.6)	(0.1)	(100.0)				
HD(F)単独.PDカテあり		1	2	10	47	42	41	10	2	3	158	24	182	150.65	25.22
(%)		(0.6)	(1.3)	(6.3)	(29.7)	(26.6)	(25.9)	(6.3)	(1.3)	(1.9)	(100.0)				
PD.HD併用週1		4	25	90	296	391	256	87	24	7	1,180	290	1,470	148.54	24.26
HD(F)															
(%)		(0.3)	(2.1)	(7.6)	(25.1)	(33.1)	(21.7)	(7.4)	(2.0)	(0.6)	(100.0)				
PD.HD併用週2			5	14	37	54	22	13	3		148	24	172	146.41	24.23
HD(F)															
(%)			(3.4)	(9.5)	(25.0)	(36.5)	(14.9)	(8.8)	(2.0)		(100.0)				
PD.HD併用週3				2	4	6	7	3	2		24	5	29	158.08	26.36
HD(F)															
(%)				(8.3)	(16.7)	(25.0)	(29.2)	(12.5)	(8.3)		(100.0)				
PD.HD併用週4															
HD(F)															
(%)															
PD.HD併用その他				7	20	18	13	3			61	10	71	144.36	21.74
(%)				(11.5)	(32.8)	(29.5)	(21.3)	(4.9)			(100.0)				
合計	46	445	3,877	19,897	59,137	89,626	66,459	25,902	6,596	1,389	273,374	39,817	313,191	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				
不明			1		5	5	1				12	13	25	136.83	17.71
(%)			(8.3)		(41.7)	(41.7)	(8.3)				(100.0)				
記載なし												1	1		
(%)															
総計	46	445	3,878	19,897	59,142	89,631	66,460	25,902	6,596	1,389	273,386	39,831	313,217	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査