

表 3246 透析前収縮期血圧(mmHg) 性別（透析患者全体）

性別	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
男性	21	238	2,151	11,752	37,367	58,442	43,850	16,822	4,210	849	175,702	25,635	201,337	152.09	24.01
(%)	(0.0)	(0.1)	(1.2)	(6.7)	(21.3)	(33.3)	(25.0)	(9.6)	(2.4)	(0.5)	(100.0)				
女性	25	207	1,727	8,145	21,775	31,189	22,610	9,080	2,386	540	97,684	14,196	111,880	150.45	25.23
(%)	(0.0)	(0.2)	(1.8)	(8.3)	(22.3)	(31.9)	(23.1)	(9.3)	(2.4)	(0.6)	(100.0)				
合計	46	445	3,878	19,897	59,142	89,631	66,460	25,902	6,596	1,389	273,386	39,831	313,217	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				
記載なし															
(%)															
総計	46	445	3,878	19,897	59,142	89,631	66,460	25,902	6,596	1,389	273,386	39,831	313,217	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査