

表 3250 透析前収縮期血圧(mmHg) 医療機関別（透析患者全体）

医療機関	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
国公立大学		4	32	104	204	258	126	40	5		773	435	1,208	141.07	24.18
(%)		(0.5)	(4.1)	(13.5)	(26.4)	(33.4)	(16.3)	(5.2)	(0.6)		(100.0)				
私立大学		8	67	219	589	823	501	187	47	10	2,451	1,113	3,564	147.71	25.10
(%)		(0.3)	(2.7)	(8.9)	(24.0)	(33.6)	(20.4)	(7.6)	(1.9)	(0.4)	(100.0)				
国立		1	21	56	145	205	147	49	11	1	636	288	924	147.74	24.78
(%)		(0.2)	(3.3)	(8.8)	(22.8)	(32.2)	(23.1)	(7.7)	(1.7)	(0.2)	(100.0)				
県市町村立	1	33	231	1,289	3,695	5,271	3,595	1,399	360	89	15,963	3,555	19,518	150.20	24.65
(%)	(0.0)	(0.2)	(1.4)	(8.1)	(23.1)	(33.0)	(22.5)	(8.8)	(2.3)	(0.6)	(100.0)				
社会保険		8	45	237	706	1,124	834	307	79	11	3,351	548	3,899	151.66	24.06
(%)		(0.2)	(1.3)	(7.1)	(21.1)	(33.5)	(24.9)	(9.2)	(2.4)	(0.3)	(100.0)				
厚生連		15	119	629	1,703	2,593	1,884	774	224	50	7,991	1,010	9,001	151.68	25.18
(%)		(0.2)	(1.5)	(7.9)	(21.3)	(32.4)	(23.6)	(9.7)	(2.8)	(0.6)	(100.0)				
その他公的	1	21	169	680	1,943	2,808	1,980	720	195	45	8,562	2,362	10,924	149.87	24.88
(%)	(0.0)	(0.2)	(2.0)	(7.9)	(22.7)	(32.8)	(23.1)	(8.4)	(2.3)	(0.5)	(100.0)				
私立総合	2	12	90	490	1,538	2,000	1,498	597	163	30	6,420	1,028	7,448	150.81	24.76
(%)	(0.0)	(0.2)	(1.4)	(7.6)	(24.0)	(31.2)	(23.3)	(9.3)	(2.5)	(0.5)	(100.0)				
私立	16	136	1,356	6,447	17,557	25,346	18,624	7,237	1,771	368	78,858	11,207	90,065	150.45	24.79
(%)	(0.0)	(0.2)	(1.7)	(8.2)	(22.3)	(32.1)	(23.6)	(9.2)	(2.2)	(0.5)	(100.0)				
私立診療所	26	207	1,748	9,746	31,062	49,203	37,271	14,592	3,741	785	148,381	18,285	166,666	152.45	24.14
(%)	(0.0)	(0.1)	(1.2)	(6.6)	(20.9)	(33.2)	(25.1)	(9.8)	(2.5)	(0.5)	(100.0)				
総計	46	445	3,878	19,897	59,142	89,631	66,460	25,902	6,596	1,389	273,386	39,831	313,217	151.51	24.47
(%)	(0.0)	(0.2)	(1.4)	(7.3)	(21.6)	(32.8)	(24.3)	(9.5)	(2.4)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査