

表 3258 透析前収縮期血圧(mmHg) 医療機関別（施設血液透析患者 透析回数週3回）

医療機関	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
国公立大学		1	16	51	77	131	78	28	3		385	23	408	144.30	25.55
(%)		(0.3)	(4.2)	(13.2)	(20.0)	(34.0)	(20.3)	(7.3)	(0.8)		(100.0)				
私立大学		5	39	118	313	465	356	134	36	8	1,474	22	1,496	150.56	25.88
(%)		(0.3)	(2.6)	(8.0)	(21.2)	(31.5)	(24.2)	(9.1)	(2.4)	(0.5)	(100.0)				
国立		1	8	36	86	147	111	39	8	1	437	78	515	150.49	24.02
(%)		(0.2)	(1.8)	(8.2)	(19.7)	(33.6)	(25.4)	(8.9)	(1.8)	(0.2)	(100.0)				
県市町村立	1	18	138	910	2,722	4,011	2,793	1,101	280	61	12,035	935	12,970	150.99	24.27
(%)	(0.0)	(0.1)	(1.1)	(7.6)	(22.6)	(33.3)	(23.2)	(9.1)	(2.3)	(0.5)	(100.0)				
社会保険		4	25	168	524	877	666	247	66	6	2,583	92	2,675	152.77	23.49
(%)		(0.2)	(1.0)	(6.5)	(20.3)	(34.0)	(25.8)	(9.6)	(2.6)	(0.2)	(100.0)				
厚生連		8	79	452	1,267	2,016	1,495	616	176	42	6,151	272	6,423	152.60	25.01
(%)		(0.1)	(1.3)	(7.3)	(20.6)	(32.8)	(24.3)	(10.0)	(2.9)	(0.7)	(100.0)				
その他公的		12	105	481	1,434	2,174	1,586	581	163	37	6,573	343	6,916	151.11	24.61
(%)		(0.2)	(1.6)	(7.3)	(21.8)	(33.1)	(24.1)	(8.8)	(2.5)	(0.6)	(100.0)				
私立総合	2	8	58	358	1,071	1,419	1,096	432	130	23	4,597	192	4,789	151.35	24.92
(%)	(0.0)	(0.2)	(1.3)	(7.8)	(23.3)	(30.9)	(23.8)	(9.4)	(2.8)	(0.5)	(100.0)				
私立	11	113	982	5,001	14,017	20,318	14,998	5,775	1,438	289	62,942	3,655	66,597	150.70	24.62
(%)	(0.0)	(0.2)	(1.6)	(7.9)	(22.3)	(32.3)	(23.8)	(9.2)	(2.3)	(0.5)	(100.0)				
私立診療所	18	143	1,208	7,158	23,509	37,839	28,816	11,274	2,843	584	113,392	7,671	121,063	152.72	23.92
(%)	(0.0)	(0.1)	(1.1)	(6.3)	(20.7)	(33.4)	(25.4)	(9.9)	(2.5)	(0.5)	(100.0)				
総計	32	313	2,658	14,733	45,020	69,397	51,995	20,227	5,143	1,051	210,569	13,283	223,852	151.90	24.26
(%)	(0.0)	(0.1)	(1.3)	(7.0)	(21.4)	(33.0)	(24.7)	(9.6)	(2.4)	(0.5)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査