

表 3266 透析前収縮期血圧(mmHg) 医療機関別（血液透析濾過患者 透析回数週3回）

医療機関	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
国公立大学			8	3	11	14	7	3	1		47	2	49	136.30	30.72
(%)			(17.0)	(6.4)	(23.4)	(29.8)	(14.9)	(6.4)	(2.1)		(100.0)				
私立大学			4	13	31	73	31	19	4		175		175	151.14	24.05
(%)			(2.3)	(7.4)	(17.7)	(41.7)	(17.7)	(10.9)	(2.3)		(100.0)				
国立			2	4	5	15	20	4			50	1	51	152.40	23.19
(%)			(4.0)	(8.0)	(10.0)	(30.0)	(40.0)	(8.0)			(100.0)				
県市町村立		5	44	163	398	620	429	179	59	18	1,915	200	2,115	151.15	26.48
(%)		(0.3)	(2.3)	(8.5)	(20.8)	(32.4)	(22.4)	(9.3)	(3.1)	(0.9)	(100.0)				
社会保険		4	12	33	89	118	90	38	11	3	398	15	413	149.43	27.48
(%)		(1.0)	(3.0)	(8.3)	(22.4)	(29.6)	(22.6)	(9.5)	(2.8)	(0.8)	(100.0)				
厚生連		2	22	85	200	279	241	111	34	6	980	42	1,022	152.25	26.52
(%)		(0.2)	(2.2)	(8.7)	(20.4)	(28.5)	(24.6)	(11.3)	(3.5)	(0.6)	(100.0)				
その他公的		4	21	63	155	246	179	77	18	5	768	86	854	150.60	26.42
(%)		(0.5)	(2.7)	(8.2)	(20.2)	(32.0)	(23.3)	(10.0)	(2.3)	(0.7)	(100.0)				
私立総合		2	19	77	259	393	296	130	22	5	1,203	39	1,242	151.79	24.15
(%)		(0.2)	(1.6)	(6.4)	(21.5)	(32.7)	(24.6)	(10.8)	(1.8)	(0.4)	(100.0)				
私立	5	18	257	959	2,412	3,643	2,758	1,163	271	62	11,548	706	12,254	150.88	25.61
(%)	(0.0)	(0.2)	(2.2)	(8.3)	(20.9)	(31.5)	(23.9)	(10.1)	(2.3)	(0.5)	(100.0)				
私立診療所	6	56	450	2,113	6,314	9,693	7,318	2,920	812	172	29,854	1,552	31,406	152.14	24.84
(%)	(0.0)	(0.2)	(1.5)	(7.1)	(21.1)	(32.5)	(24.5)	(9.8)	(2.7)	(0.6)	(100.0)				
総計	11	91	839	3,513	9,874	15,094	11,369	4,644	1,232	271	46,938	2,643	49,581	151.71	25.18
(%)	(0.0)	(0.2)	(1.8)	(7.5)	(21.0)	(32.2)	(24.2)	(9.9)	(2.6)	(0.6)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査