

表 3274 収縮期血圧(mmHg) 医療機関別（腹膜透析患者）

医療機関	60未満	60～	80～	100～	120～	140～	160～	180～	200～	220～	合計	記載なし	総計	平均	標準偏差
国公立大学		3	7	45	108	106	31	8	1		309	354	663	137.56	20.53
(%)		(1.0)	(2.3)	(14.6)	(35.0)	(34.3)	(10.0)	(2.6)	(0.3)		(100.0)				
私立大学		2	19	64	191	226	82	24	5	1	614	492	1,106	141.40	21.91
(%)		(0.3)	(3.1)	(10.4)	(31.1)	(36.8)	(13.4)	(3.9)	(0.8)	(0.2)	(100.0)				
国立			9	12	42	37	14	3	3		120	102	222	138.13	24.87
(%)			(7.5)	(10.0)	(35.0)	(30.8)	(11.7)	(2.5)	(2.5)		(100.0)				
県市町村立		6	33	124	355	298	132	43	5	1	997	576	1,573	139.42	22.79
(%)		(0.6)	(3.3)	(12.4)	(35.6)	(29.9)	(13.2)	(4.3)	(0.5)	(0.1)	(100.0)				
社会保険			4	19	48	65	34	4	1	1	176	346	522	143.26	21.90
(%)			(2.3)	(10.8)	(27.3)	(36.9)	(19.3)	(2.3)	(0.6)	(0.6)	(100.0)				
厚生連		3	11	45	129	150	60	15	4	1	418	142	560	141.68	22.32
(%)		(0.7)	(2.6)	(10.8)	(30.9)	(35.9)	(14.4)	(3.6)	(1.0)	(0.2)	(100.0)				
その他公的		4	33	93	269	275	146	36	9		865	494	1,359	141.56	23.35
(%)		(0.5)	(3.8)	(10.8)	(31.1)	(31.8)	(16.9)	(4.2)	(1.0)		(100.0)				
私立総合		1	8	31	111	93	37	11	2	2	296	67	363	141.06	22.60
(%)		(0.3)	(2.7)	(10.5)	(37.5)	(31.4)	(12.5)	(3.7)	(0.7)	(0.7)	(100.0)				
私立			39	133	325	291	160	31	12	4	995	704	1,699	140.66	23.55
(%)			(3.9)	(13.4)	(32.7)	(29.2)	(16.1)	(3.1)	(1.2)	(0.4)	(100.0)				
私立診療所		1	6	54	145	173	104	34	5	1	523	179	702	145.89	22.60
(%)		(0.2)	(1.1)	(10.3)	(27.7)	(33.1)	(19.9)	(6.5)	(1.0)	(0.2)	(100.0)				
総計		20	169	620	1,723	1,714	800	209	47	11	5,313	3,456	8,769	141.13	22.83
(%)		(0.4)	(3.2)	(11.7)	(32.4)	(32.3)	(15.1)	(3.9)	(0.9)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査