

表 3279 透析前拡張期血圧(mmHg) 治療方法(新分類)別(透析患者全体)

治療方法 (新分類)	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
施設血液透析	76	688	18,943	101,499	53,056	28,740	14,097	922	66	218,087	30,638	248,725	77.64	14.46
(%)	(0.0)	(0.3)	(8.7)	(46.5)	(24.3)	(13.2)	(6.5)	(0.4)	(0.0)	(100.0)				
Off-line HDF	4	30	497	2,101	1,105	601	316	21	1	4,676	656	5,332	76.77	15.40
(%)	(0.1)	(0.6)	(10.6)	(44.9)	(23.6)	(12.9)	(6.8)	(0.4)	(0.0)	(100.0)				
On-line HDF	14	114	2,731	16,446	10,303	6,277	3,479	237	13	39,614	4,913	44,527	80.05	14.73
(%)	(0.0)	(0.3)	(6.9)	(41.5)	(26.0)	(15.8)	(8.8)	(0.6)	(0.0)	(100.0)				
Push/Pull HDF		1	7	45	28	15	5			101	9	110	77.83	14.06
(%)		(1.0)	(6.9)	(44.6)	(27.7)	(14.9)	(5.0)			(100.0)				
ハイオフィルトレーション		2	49	103	46	24	16	2		242	25	267	73.78	16.78
(%)		(0.8)	(20.2)	(42.6)	(19.0)	(9.9)	(6.6)	(0.8)		(100.0)				
IHDF		20	346	1,531	790	431	204	11	1	3,334	206	3,540	77.17	14.78
(%)		(0.6)	(10.4)	(45.9)	(23.7)	(12.9)	(6.1)	(0.3)	(0.0)	(100.0)				
血液濾過			2	5	3	1				11	5	16	74.64	15.52
(%)			(18.2)	(45.5)	(27.3)	(9.1)				(100.0)				
血液吸着透析 (リクセル等使用)	1	21	165	619	267	113	51	4	2	1,243	133	1,376	73.93	15.40
(%)	(0.1)	(1.7)	(13.3)	(49.8)	(21.5)	(9.1)	(4.1)	(0.3)	(0.2)	(100.0)				
在宅血液透析			21	144	98	58	33	4		358	197	555	80.69	14.41
(%)			(5.9)	(40.2)	(27.4)	(16.2)	(9.2)	(1.1)		(100.0)				
腹膜透析 (CAPD)		4	209	1,303	721	432	212	18	1	2,900	2,002	4,902	78.78	14.59
(%)		(0.1)	(7.2)	(44.9)	(24.9)	(14.9)	(7.3)	(0.6)	(0.0)	(100.0)				
腹膜透析 (APD)	3	1	136	667	447	282	120	3		1,659	1,116	2,775	79.00	14.63
(%)	(0.2)	(0.1)	(8.2)	(40.2)	(26.9)	(17.0)	(7.2)	(0.2)		(100.0)				
腹膜透析 (CCPD)		1	35	277	210	106	86	5		720	372	1,092	81.68	14.47
(%)		(0.1)	(4.9)	(38.5)	(29.2)	(14.7)	(11.9)	(0.7)		(100.0)				
合計	98	882	23,141	124,740	67,074	37,080	18,619	1,227	84	272,945	40,272	313,217	77.99	14.56
(%)	(0.0)	(0.3)	(8.5)	(45.7)	(24.6)	(13.6)	(6.8)	(0.4)	(0.0)	(100.0)				
記載なし														
(%)														
総計	98	882	23,141	124,740	67,074	37,080	18,619	1,227	84	272,945	40,272	313,217	77.99	14.56
(%)	(0.0)	(0.3)	(8.5)	(45.7)	(24.6)	(13.6)	(6.8)	(0.4)	(0.0)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

*腹膜透析(CCPD)は、APDと手動のバック交換の両方を行うもの

(C)Japanese Society for Dialysis Therapy

2015年調査