

表 3284 透析前拡張期血圧(mmHg) 原疾患別（透析患者全体）

原疾患	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	30	296	6,248	36,023	20,895	11,835	5,614	358	23	81,322	12,025	93,347	78.56	14.43
(%)	(0.0)	(0.4)	(7.7)	(44.3)	(25.7)	(14.6)	(6.9)	(0.4)	(0.0)	(100.0)				
慢性腎盂腎炎	2	14	225	1,197	651	332	181	10		2,612	323	2,935	77.70	14.58
(%)	(0.1)	(0.5)	(8.6)	(45.8)	(24.9)	(12.7)	(6.9)	(0.4)		(100.0)				
急速進行性糸球体腎炎	1		178	1,050	506	276	134	8		2,153	325	2,478	77.72	14.02
(%)	(0.0)		(8.3)	(48.8)	(23.5)	(12.8)	(6.2)	(0.4)		(100.0)				
妊娠腎/妊娠中毒症	1	6	99	607	371	206	92	4		1,386	159	1,545	78.58	14.36
(%)	(0.1)	(0.4)	(7.1)	(43.8)	(26.8)	(14.9)	(6.6)	(0.3)		(100.0)				
その他の分類不能の腎炎		3	91	520	309	191	104	5		1,223	182	1,405	79.63	14.59
(%)		(0.2)	(7.4)	(42.5)	(25.3)	(15.6)	(8.5)	(0.4)		(100.0)				
多発性嚢胞腎	1	20	516	4,256	2,800	1,595	699	18		9,905	1,351	11,256	79.91	13.36
(%)	(0.0)	(0.2)	(5.2)	(43.0)	(28.3)	(16.1)	(7.1)	(0.2)		(100.0)				
腎硬化症	10	97	2,758	12,797	5,860	3,002	1,438	101	11	26,074	3,731	29,805	76.17	14.50
(%)	(0.0)	(0.4)	(10.6)	(49.1)	(22.5)	(11.5)	(5.5)	(0.4)	(0.0)	(100.0)				
悪性高血圧	1	5	140	857	565	387	293	18	1	2,267	351	2,618	81.83	15.60
(%)	(0.0)	(0.2)	(6.2)	(37.8)	(24.9)	(17.1)	(12.9)	(0.8)	(0.0)	(100.0)				
糖尿病性腎症	32	276	9,061	48,874	25,318	13,617	7,239	476	42	104,935	15,343	120,278	77.80	14.55
(%)	(0.0)	(0.3)	(8.6)	(46.6)	(24.1)	(13.0)	(6.9)	(0.5)	(0.0)	(100.0)				
SLE腎炎		7	140	806	471	304	183	13	1	1,925	302	2,227	79.68	15.16
(%)		(0.4)	(7.3)	(41.9)	(24.5)	(15.8)	(9.5)	(0.7)	(0.1)	(100.0)				
アミロイド腎		3	30	179	97	59	18	2		388	74	462	77.40	14.09
(%)		(0.8)	(7.7)	(46.1)	(25.0)	(15.2)	(4.6)	(0.5)		(100.0)				
痛風腎	2	8	73	421	265	113	65	2		949	126	1,075	77.88	14.45
(%)	(0.2)	(0.8)	(7.7)	(44.4)	(27.9)	(11.9)	(6.8)	(0.2)		(100.0)				
先天性代謝異常による腎不全	1	5	19	87	61	36	16	5		230	44	274	79.29	17.58
(%)	(0.4)	(2.2)	(8.3)	(37.8)	(26.5)	(15.7)	(7.0)	(2.2)		(100.0)				
腎・尿路結核		1	21	89	37	18	4			170	25	195	73.91	12.72
(%)		(0.6)	(12.4)	(52.4)	(21.8)	(10.6)	(2.4)			(100.0)				
腎・尿路結石		3	57	247	119	57	33			516	65	581	76.15	14.42
(%)		(0.6)	(11.0)	(47.9)	(23.1)	(11.0)	(6.4)			(100.0)				
腎・尿路腫瘍			71	385	208	96	37	3		800	110	910	76.94	13.47
(%)			(8.9)	(48.1)	(26.0)	(12.0)	(4.6)	(0.4)		(100.0)				
閉塞性尿路障害		2	48	274	149	106	48	9		636	94	730	79.30	15.20
(%)		(0.3)	(7.5)	(43.1)	(23.4)	(16.7)	(7.5)	(1.4)		(100.0)				
骨髄腫		2	34	125	58	26	4			249	42	291	73.55	13.40
(%)		(0.8)	(13.7)	(50.2)	(23.3)	(10.4)	(1.6)			(100.0)				
腎形成不全			45	211	145	106	63	10		580	80	660	81.76	15.89
(%)			(7.8)	(36.4)	(25.0)	(18.3)	(10.9)	(1.7)		(100.0)				
不明	13	91	2,463	11,860	6,119	3,419	1,699	131	4	25,799	4,098	29,897	77.48	14.85
(%)	(0.1)	(0.4)	(9.5)	(46.0)	(23.7)	(13.3)	(6.6)	(0.5)	(0.0)	(100.0)				
再導入		10	113	689	506	356	191	12		1,877	306	2,183	81.28	15.13
(%)		(0.5)	(6.0)	(36.7)	(27.0)	(19.0)	(10.2)	(0.6)		(100.0)				
その他	4	33	709	3,180	1,561	941	462	42	2	6,934	1,110	8,044	77.25	15.13
(%)	(0.1)	(0.5)	(10.2)	(45.9)	(22.5)	(13.6)	(6.7)	(0.6)	(0.0)	(100.0)				
合計	98	882	23,139	124,734	67,071	37,078	18,617	1,227	84	272,930	40,266	313,196	77.99	14.56
(%)	(0.0)	(0.3)	(8.5)	(45.7)	(24.6)	(13.6)	(6.8)	(0.4)	(0.0)	(100.0)				
記載なし			2	6	3	2	2			15	6	21	79.53	17.27
(%)			(13.3)	(40.0)	(20.0)	(13.3)	(13.3)			(100.0)				
総計	98	882	23,141	124,740	67,074	37,080	18,619	1,227	84	272,945	40,272	313,217	77.99	14.56
(%)	(0.0)	(0.3)	(8.5)	(45.7)	(24.6)	(13.6)	(6.8)	(0.4)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査