

表 3291 透析前拡張期血圧(mmHg) 原疾患別（施設血液透析患者 透析回数週3回）

原疾患	20未満	20～	40～	60～	80～	90～	100～	120～	140～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	22	201	4,640	26,871	15,252	8,502	3,931	250	17	59,686	3,788	63,474	78.30	14.32
(%)	(0.0)	(0.3)	(7.8)	(45.0)	(25.6)	(14.2)	(6.6)	(0.4)	(0.0)	(100.0)				
慢性腎盂腎炎	1	10	166	902	495	227	128	9		1,938	102	2,040	77.46	14.39
(%)	(0.1)	(0.5)	(8.6)	(46.5)	(25.5)	(11.7)	(6.6)	(0.5)		(100.0)				
急速進行性糸球体腎炎	1		140	818	389	222	102	5		1,677	94	1,771	77.64	14.00
(%)	(0.1)		(8.3)	(48.8)	(23.2)	(13.2)	(6.1)	(0.3)		(100.0)				
妊娠腎/妊娠中毒症		3	68	415	242	139	56	1		924	57	981	78.18	13.92
(%)		(0.3)	(7.4)	(44.9)	(26.2)	(15.0)	(6.1)	(0.1)		(100.0)				
その他の分類不能の腎炎		1	67	384	213	130	70	5		870	59	929	79.17	14.68
(%)		(0.1)	(7.7)	(44.1)	(24.5)	(14.9)	(8.0)	(0.6)		(100.0)				
多発性嚢胞腎	1	16	382	3,294	2,111	1,199	512	13		7,528	488	8,016	79.75	13.32
(%)	(0.0)	(0.2)	(5.1)	(43.8)	(28.0)	(15.9)	(6.8)	(0.2)		(100.0)				
腎硬化症	8	76	2,191	10,392	4,627	2,328	1,094	75	9	20,800	1,207	22,007	75.95	14.35
(%)	(0.0)	(0.4)	(10.5)	(50.0)	(22.2)	(11.2)	(5.3)	(0.4)	(0.0)	(100.0)				
悪性高血圧	1	3	114	700	409	294	217	13	1	1,752	109	1,861	81.13	15.69
(%)	(0.1)	(0.2)	(6.5)	(40.0)	(23.3)	(16.8)	(12.4)	(0.7)	(0.1)	(100.0)				
糖尿病性腎症	26	217	7,409	39,582	20,108	10,636	5,568	359	31	83,936	5,490	89,426	77.54	14.47
(%)	(0.0)	(0.3)	(8.8)	(47.2)	(24.0)	(12.7)	(6.6)	(0.4)	(0.0)	(100.0)				
SLE腎炎		4	99	590	343	229	127	8	1	1,401	103	1,504	79.57	14.92
(%)		(0.3)	(7.1)	(42.1)	(24.5)	(16.3)	(9.1)	(0.6)	(0.1)	(100.0)				
アミロイド腎	1	18	138	138	73	46	12	1		289	25	314	77.68	13.55
(%)		(0.3)	(6.2)	(47.8)	(25.3)	(15.9)	(4.2)	(0.3)		(100.0)				
痛風腎	2	7	61	327	206	87	48	1		739	39	778	77.50	14.59
(%)	(0.3)	(0.9)	(8.3)	(44.2)	(27.9)	(11.8)	(6.5)	(0.1)		(100.0)				
先天性代謝異常による腎不全	1	5	13	66	42	22	10	5		164	9	173	78.15	18.50
(%)	(0.6)	(3.0)	(7.9)	(40.2)	(25.6)	(13.4)	(6.1)	(3.0)		(100.0)				
腎・尿路結核		1	17	67	30	15	3			133	8	141	73.58	13.22
(%)		(0.8)	(12.8)	(50.4)	(22.6)	(11.3)	(2.3)			(100.0)				
腎・尿路結石		3	44	199	87	44	26			403	18	421	75.81	14.33
(%)		(0.7)	(10.9)	(49.4)	(21.6)	(10.9)	(6.5)			(100.0)				
腎・尿路腫瘍			60	310	169	76	30	2		647	31	678	76.89	13.43
(%)			(9.3)	(47.9)	(26.1)	(11.7)	(4.6)	(0.3)		(100.0)				
閉塞性尿路障害		2	32	206	110	77	35	8		470	24	494	79.37	15.15
(%)		(0.4)	(6.8)	(43.8)	(23.4)	(16.4)	(7.4)	(1.7)		(100.0)				
骨髄腫	1	18	98	43	17	3				180	10	190	74.24	12.36
(%)		(0.6)	(10.0)	(54.4)	(23.9)	(9.4)	(1.7)			(100.0)				
腎形成不全			26	136	102	69	43	7		383	17	400	82.29	15.53
(%)			(6.8)	(35.5)	(26.6)	(18.0)	(11.2)	(1.8)		(100.0)				
不明	9	68	1,966	9,210	4,605	2,533	1,228	93	4	19,716	1,460	21,176	77.07	14.79
(%)	(0.0)	(0.3)	(10.0)	(46.7)	(23.4)	(12.8)	(6.2)	(0.5)	(0.0)	(100.0)				
再導入		7	75	473	367	236	128	10		1,296	60	1,356	81.37	14.99
(%)		(0.5)	(5.8)	(36.5)	(28.3)	(18.2)	(9.9)	(0.8)		(100.0)				
その他	4	27	525	2,497	1,182	710	341	32	2	5,320	395	5,715	77.03	15.02
(%)	(0.1)	(0.5)	(9.9)	(46.9)	(22.2)	(13.3)	(6.4)	(0.6)	(0.0)	(100.0)				
合計	76	653	18,131	97,675	51,205	27,838	13,712	897	65	210,252	13,593	223,845	77.70	14.46
(%)	(0.0)	(0.3)	(8.6)	(46.5)	(24.4)	(13.2)	(6.5)	(0.4)	(0.0)	(100.0)				
記載なし			2	4						6	1	7	65.33	11.41
(%)			(33.3)	(66.7)						(100.0)				
総計	76	653	18,133	97,679	51,205	27,838	13,712	897	65	210,258	13,594	223,852	77.70	14.46
(%)	(0.0)	(0.3)	(8.6)	(46.5)	(24.4)	(13.2)	(6.5)	(0.4)	(0.0)	(100.0)				

数値下のかつこ内は行方向の合計に対する％です。
(C)Japanese Society for Dialysis Therapy
2015年調査