

表 3318 透析前脈拍(拍/分) 性別（施設血液透析患者 透析回数週3回）

性別	40未満	40～	50～	60～	70～	80～	90～	100～	110～	120～	合計	記載なし	総計	平均	標準偏差
男性	108	1,665	12,692	35,736	40,178	25,542	11,172	3,474	831	313	131,711	12,729	144,440	74.05	12.91
(%)	(0.1)	(1.3)	(9.6)	(27.1)	(30.5)	(19.4)	(8.5)	(2.6)	(0.6)	(0.2)	(100.0)				
女性	60	592	5,322	19,200	24,233	14,851	5,896	1,767	404	179	72,504	6,908	79,412	74.67	12.29
(%)	(0.1)	(0.8)	(7.3)	(26.5)	(33.4)	(20.5)	(8.1)	(2.4)	(0.6)	(0.2)	(100.0)				
合計	168	2,257	18,014	54,936	64,411	40,393	17,068	5,241	1,235	492	204,215	19,637	223,852	74.27	12.70
(%)	(0.1)	(1.1)	(8.8)	(26.9)	(31.5)	(19.8)	(8.4)	(2.6)	(0.6)	(0.2)	(100.0)				
記載なし															
(%)															
総計	168	2,257	18,014	54,936	64,411	40,393	17,068	5,241	1,235	492	204,215	19,637	223,852	74.27	12.70
(%)	(0.1)	(1.1)	(8.8)	(26.9)	(31.5)	(19.8)	(8.4)	(2.6)	(0.6)	(0.2)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。
(C)Japanese Society for Dialysis Therapy
2015年調査