

表 3404 一日透析液使用量(L/日) 原疾患別(腹膜透析単独患者)

原疾患	2未満	2～	4～	6～	8～	10～	12～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	32	82	298	528	353	81	36	1,410	736	2,146	6.55	2.31
(%)	(2.3)	(5.8)	(21.1)	(37.4)	(25.0)	(5.7)	(2.6)	(100.0)				
慢性腎盂腎炎	3	5	11	24	10	2	2	57	21	78	6.23	2.65
(%)	(5.3)	(8.8)	(19.3)	(42.1)	(17.5)	(3.5)	(3.5)	(100.0)				
急速進行性糸球体腎炎	1	3	3	11	9	4	2	33	22	55	7.23	2.90
(%)	(3.0)	(9.1)	(9.1)	(33.3)	(27.3)	(12.1)	(6.1)	(100.0)				
妊娠腎/妊娠中毒症			4	4	4	2		14	4	18	6.92	2.16
(%)			(28.6)	(28.6)	(28.6)	(14.3)		(100.0)				
その他の分類不能の腎炎		2	11	8	9	1	1	32	27	59	6.69	2.62
(%)		(6.3)	(34.4)	(25.0)	(28.1)	(3.1)	(3.1)	(100.0)				
多発性嚢胞腎	8	18	35	56	20	8		145	80	225	5.72	2.24
(%)	(5.5)	(12.4)	(24.1)	(38.6)	(13.8)	(5.5)		(100.0)				
腎硬化症	15	80	144	225	124	34	10	632	311	943	6.12	2.35
(%)	(2.4)	(12.7)	(22.8)	(35.6)	(19.6)	(5.4)	(1.6)	(100.0)				
悪性高血圧	7	6	14	12	15		1	55	32	87	5.53	2.57
(%)	(12.7)	(10.9)	(25.5)	(21.8)	(27.3)		(1.8)	(100.0)				
糖尿病性腎症	33	83	302	559	329	94	24	1,424	902	2,326	6.55	2.31
(%)	(2.3)	(5.8)	(21.2)	(39.3)	(23.1)	(6.6)	(1.7)	(100.0)				
SLE腎炎	3	1	6	5	4	4		23	17	40	6.09	3.26
(%)	(13.0)	(4.3)	(26.1)	(21.7)	(17.4)	(17.4)		(100.0)				
アミロイド腎		2	2	2	1			7	6	13	5.16	1.80
(%)		(28.6)	(28.6)	(28.6)	(14.3)			(100.0)				
痛風腎			3	5	7	2	1	18	5	23	7.80	2.20
(%)			(16.7)	(27.8)	(38.9)	(11.1)	(5.6)	(100.0)				
先天性代謝異常による腎不全	1		3	4	2		1	11	4	15	6.46	3.41
(%)	(9.1)		(27.3)	(36.4)	(18.2)		(9.1)	(100.0)				
腎・尿路結核		1		1				2		2	4.75	1.77
(%)		(50.0)		(50.0)				(100.0)				
腎・尿路結石	1	1	1	6	3	1	1	14	3	17	6.72	2.73
(%)	(7.1)	(7.1)	(7.1)	(42.9)	(21.4)	(7.1)	(7.1)	(100.0)				
腎・尿路腫瘍			1	2	2	1		6	4	10	7.50	1.83
(%)			(16.7)	(33.3)	(33.3)	(16.7)		(100.0)				
閉塞性尿路障害		3	4	4	5	1		17	2	19	6.31	2.13
(%)		(17.6)	(23.5)	(23.5)	(29.4)	(5.9)		(100.0)				
骨髄腫			2	1	2			5	2	7	6.70	2.17
(%)			(40.0)	(20.0)	(40.0)			(100.0)				
腎形成不全	4	8	9	6	9	1	7	44	23	67	7.45	5.69
(%)	(9.1)	(18.2)	(20.5)	(13.6)	(20.5)	(2.3)	(15.9)	(100.0)				
不明	13	48	107	148	102	14	9	441	241	682	6.13	2.45
(%)	(2.9)	(10.9)	(24.3)	(33.6)	(23.1)	(3.2)	(2.0)	(100.0)				
再導入	1	1	2	10	8		2	24	17	41	7.68	4.92
(%)	(4.2)	(4.2)	(8.3)	(41.7)	(33.3)		(8.3)	(100.0)				
その他	6	11	35	53	22	7	13	147	84	231	6.86	3.97
(%)	(4.1)	(7.5)	(23.8)	(36.1)	(15.0)	(4.8)	(8.8)	(100.0)				
合計	128	355	997	1,674	1,040	257	110	4,561	2,543	7,104	6.44	2.51
(%)	(2.8)	(7.8)	(21.9)	(36.7)	(22.8)	(5.6)	(2.4)	(100.0)				
記載なし												
(%)												
総計	128	355	997	1,674	1,040	257	110	4,561	2,543	7,104	6.44	2.51
(%)	(2.8)	(7.8)	(21.9)	(36.7)	(22.8)	(5.6)	(2.4)	(100.0)				

数値下のカッコ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査