

表 3416 残存腎機能(一日尿量)(mL/日) 年齢(歳)別(腹膜透析患者)

年齢	100未満	100～	400～	800～	1200～	1600～	合計	記載なし	総計	平均	標準偏差
15歳未満	29	7	11	2		2	51	38	89	275.47	537.24
(%)	(56.9)	(13.7)	(21.6)	(3.9)		(3.9)	(100.0)				
15歳～	25	6	9	8	2	2	52	49	101	397.15	546.24
(%)	(48.1)	(11.5)	(17.3)	(15.4)	(3.8)	(3.8)	(100.0)				
30歳～	79	63	65	60	30	30	327	333	660	649.46	655.57
(%)	(24.2)	(19.3)	(19.9)	(18.3)	(9.2)	(9.2)	(100.0)				
45歳～	305	210	255	259	126	85	1,240	1,080	2,320	622.73	587.38
(%)	(24.6)	(16.9)	(20.6)	(20.9)	(10.2)	(6.9)	(100.0)				
60歳～	405	404	500	436	218	137	2,100	1,723	3,823	643.60	569.13
(%)	(19.3)	(19.2)	(23.8)	(20.8)	(10.4)	(6.5)	(100.0)				
75歳～	110	165	248	200	72	43	838	811	1,649	648.46	496.92
(%)	(13.1)	(19.7)	(29.6)	(23.9)	(8.6)	(5.1)	(100.0)				
90歳～	6	15	20	8	6	2	57	70	127	582.89	492.55
(%)	(10.5)	(26.3)	(35.1)	(14.0)	(10.5)	(3.5)	(100.0)				
合計	959	870	1,108	973	454	301	4,665	4,104	8,769	631.82	568.54
(%)	(20.6)	(18.6)	(23.8)	(20.9)	(9.7)	(6.5)	(100.0)				
不明											
(%)											
記載なし											
(%)											
総計	959	870	1,108	973	454	301	4,665	4,104	8,769	631.82	568.54
(%)	(20.6)	(18.6)	(23.8)	(20.9)	(9.7)	(6.5)	(100.0)				
平均	58.42	63.65	64.82	63.76	63.40	61.16	62.69	62.93	62.80		
標準偏差	16.26	13.98	14.43	12.93	12.04	13.57	14.37	14.81	14.58		

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査