

表 3458 APD使用の有無 原疾患別（腹膜透析患者）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	1,068	775	1,843	7	980	2,830
(%)	(57.9)	(42.1)	(100.0)			
慢性腎盂腎炎	36	35	71		26	97
(%)	(50.7)	(49.3)	(100.0)			
急速進行性糸球体腎炎	16	25	41		25	66
(%)	(39.0)	(61.0)	(100.0)			
妊娠腎/妊娠中毒症	11	10	21		6	27
(%)	(52.4)	(47.6)	(100.0)			
その他の分類不能の腎炎	20	27	47		31	78
(%)	(42.6)	(57.4)	(100.0)			
多発性嚢胞腎	82	93	175		88	263
(%)	(46.9)	(53.1)	(100.0)			
腎硬化症	393	321	714	2	379	1,095
(%)	(55.0)	(45.0)	(100.0)			
悪性高血圧	34	31	65		42	107
(%)	(52.3)	(47.7)	(100.0)			
糖尿病性腎症	898	836	1,734	3	1,062	2,799
(%)	(51.8)	(48.2)	(100.0)			
SLE腎炎	17	11	28		18	46
(%)	(60.7)	(39.3)	(100.0)			
アミロイド腎	7	2	9		6	15
(%)	(77.8)	(22.2)	(100.0)			
痛風腎	10	13	23		9	32
(%)	(43.5)	(56.5)	(100.0)			
先天性代謝異常による腎不全	6	7	13		4	17
(%)	(46.2)	(53.8)	(100.0)			
腎・尿路結核	2		2			2
(%)	(100.0)		(100.0)			
腎・尿路結石	7	10	17		4	21
(%)	(41.2)	(58.8)	(100.0)			
腎・尿路腫瘍	4	5	9		5	14
(%)	(44.4)	(55.6)	(100.0)			
閉塞性尿路障害	8	8	16		5	21
(%)	(50.0)	(50.0)	(100.0)			
骨髄腫	2	2	4		3	7
(%)	(50.0)	(50.0)	(100.0)			
腎形成不全	12	36	48		24	72
(%)	(25.0)	(75.0)	(100.0)			
不明	294	233	527	1	314	842
(%)	(55.8)	(44.2)	(100.0)			
再導入	16	16	32	1	22	55
(%)	(50.0)	(50.0)	(100.0)			
その他	83	92	175		88	263
(%)	(47.4)	(52.6)	(100.0)			
合計	3,026	2,588	5,614	14	3,141	8,769
(%)	(53.9)	(46.1)	(100.0)			
記載なし						
(%)						
総計	3,026	2,588	5,614	14	3,141	8,769
(%)	(53.9)	(46.1)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査