

表 3464 APD使用の有無 原疾患別（腹膜透析単独患者）

原疾患	なし	あり	合計	不明	記載なし	総計
慢性糸球体腎炎	834	570	1,404	4	738	2,146
(%)	(59.4)	(40.6)	(100.0)			
慢性腎盂腎炎	28	30	58		20	78
(%)	(48.3)	(51.7)	(100.0)			
急速進行性糸球体腎炎	14	19	33		22	55
(%)	(42.4)	(57.6)	(100.0)			
妊娠腎/妊娠中毒症	7	7	14		4	18
(%)	(50.0)	(50.0)	(100.0)			
その他の分類不能の腎炎	17	18	35		24	59
(%)	(48.6)	(51.4)	(100.0)			
多発性嚢胞腎	70	75	145		80	225
(%)	(48.3)	(51.7)	(100.0)			
腎硬化症	348	278	626	1	316	943
(%)	(55.6)	(44.4)	(100.0)			
悪性高血圧	28	25	53		34	87
(%)	(52.8)	(47.2)	(100.0)			
糖尿病性腎症	772	680	1,452	3	871	2,326
(%)	(53.2)	(46.8)	(100.0)			
SLE腎炎	16	8	24		16	40
(%)	(66.7)	(33.3)	(100.0)			
アミロイド腎	7	2	9		4	13
(%)	(77.8)	(22.2)	(100.0)			
痛風腎	6	11	17		6	23
(%)	(35.3)	(64.7)	(100.0)			
先天性代謝異常による腎不全	6	6	12		3	15
(%)	(50.0)	(50.0)	(100.0)			
腎・尿路結核	2		2			2
(%)	(100.0)		(100.0)			
腎・尿路結石	6	8	14		3	17
(%)	(42.9)	(57.1)	(100.0)			
腎・尿路腫瘍	4	3	7		3	10
(%)	(57.1)	(42.9)	(100.0)			
閉塞性尿路障害	8	8	16		3	19
(%)	(50.0)	(50.0)	(100.0)			
骨髄腫	2	2	4		3	7
(%)	(50.0)	(50.0)	(100.0)			
腎形成不全	10	34	44		23	67
(%)	(22.7)	(77.3)	(100.0)			
不明	255	187	442	1	239	682
(%)	(57.7)	(42.3)	(100.0)			
再導入	14	10	24		17	41
(%)	(58.3)	(41.7)	(100.0)			
その他	73	80	153		78	231
(%)	(47.7)	(52.3)	(100.0)			
合計	2,527	2,061	4,588	9	2,507	7,104
(%)	(55.1)	(44.9)	(100.0)			
記載なし						
(%)						
総計	2,527	2,061	4,588	9	2,507	7,104
(%)	(55.1)	(44.9)	(100.0)			

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査