

表 3504 PD透析液交換方法 原疾患別（腹膜透析単独患者）

原疾患	完全手動式	バッグ交換デバイス(紫外線)	バッグ交換デバイス(熱式無菌接合装置)	バッグ交換デバイス(左記以外、半手動も含む)	合計	不明	記載なし	総計
慢性糸球体腎炎	479	691	199	45	1,414	4	728	2,146
(%)	(33.9)	(48.9)	(14.1)	(3.2)	(100.0)			
慢性腎盂腎炎	15	32	6	2	55	2	21	78
(%)	(27.3)	(58.2)	(10.9)	(3.6)	(100.0)			
急速進行性糸球体腎炎	2	16	10	3	31		24	55
(%)	(6.5)	(51.6)	(32.3)	(9.7)	(100.0)			
妊娠腎/妊娠中毒症	7	5	2		14		4	18
(%)	(50.0)	(35.7)	(14.3)		(100.0)			
その他の分類不能の腎炎	15	17	2		34	1	24	59
(%)	(44.1)	(50.0)	(5.9)		(100.0)			
多発性嚢胞腎	40	76	19	8	143	1	81	225
(%)	(28.0)	(53.1)	(13.3)	(5.6)	(100.0)			
腎硬化症	191	325	93	19	628	2	313	943
(%)	(30.4)	(51.8)	(14.8)	(3.0)	(100.0)			
悪性高血圧	11	34	8		53		34	87
(%)	(20.8)	(64.2)	(15.1)		(100.0)			
糖尿病性腎症	364	827	208	29	1,428	2	896	2,326
(%)	(25.5)	(57.9)	(14.6)	(2.0)	(100.0)			
SLE腎炎	11	11	2		24		16	40
(%)	(45.8)	(45.8)	(8.3)		(100.0)			
アミロイド腎	5	2	1	1	9		4	13
(%)	(55.6)	(22.2)	(11.1)	(11.1)	(100.0)			
痛風腎	8	7	3		18		5	23
(%)	(44.4)	(38.9)	(16.7)		(100.0)			
先天性代謝異常による腎不全	2	7	2		11	1	3	15
(%)	(18.2)	(63.6)	(18.2)		(100.0)			
腎・尿路結核	1	1			2			2
(%)	(50.0)	(50.0)			(100.0)			
腎・尿路結石	6	6	2		14		3	17
(%)	(42.9)	(42.9)	(14.3)		(100.0)			
腎・尿路腫瘍	3	2	1	1	7		3	10
(%)	(42.9)	(28.6)	(14.3)	(14.3)	(100.0)			
閉塞性尿路障害	2	9	5		16		3	19
(%)	(12.5)	(56.3)	(31.3)		(100.0)			
骨髄腫		4			4		3	7
(%)		(100.0)			(100.0)			
腎形成不全	16	18	8	1	43		24	67
(%)	(37.2)	(41.9)	(18.6)	(2.3)	(100.0)			
不明	144	207	67	9	427	2	253	682
(%)	(33.7)	(48.5)	(15.7)	(2.1)	(100.0)			
再導入	7	10	6		23		18	41
(%)	(30.4)	(43.5)	(26.1)		(100.0)			
その他	43	75	28	4	150	1	80	231
(%)	(28.7)	(50.0)	(18.7)	(2.7)	(100.0)			
合計	1,372	2,382	672	122	4,548	16	2,540	7,104
(%)	(30.2)	(52.4)	(14.8)	(2.7)	(100.0)			
記載なし								
(%)								
総計	1,372	2,382	672	122	4,548	16	2,540	7,104
(%)	(30.2)	(52.4)	(14.8)	(2.7)	(100.0)			

数値下のかつこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査