

表 3580 PET Cr D/P比 原疾患別（腹膜透析単独患者）

原疾患	0.5未満	0.5～	0.65～	0.81～	合計	記載なし	総計	平均	標準偏差
慢性糸球体腎炎	91	320	342	98	851	1,295	2,146	0.65	0.14
(%)	(10.7)	(37.6)	(40.2)	(11.5)	(100.0)				
慢性腎盂腎炎	7	9	11	2	29	49	78	0.60	0.15
(%)	(24.1)	(31.0)	(37.9)	(6.9)	(100.0)				
急速進行性糸球体腎炎	3	8	7	2	20	35	55	0.65	0.13
(%)	(15.0)	(40.0)	(35.0)	(10.0)	(100.0)				
妊娠腎/妊娠中毒症	1	6	3	2	12	6	18	0.66	0.12
(%)	(8.3)	(50.0)	(25.0)	(16.7)	(100.0)				
その他の分類不能の腎炎		10	8		18	41	59	0.64	0.07
(%)		(55.6)	(44.4)		(100.0)				
多発性嚢胞腎	9	36	29	8	82	143	225	0.64	0.13
(%)	(11.0)	(43.9)	(35.4)	(9.8)	(100.0)				
腎硬化症	25	94	145	49	313	630	943	0.68	0.13
(%)	(8.0)	(30.0)	(46.3)	(15.7)	(100.0)				
悪性高血圧	9	7	7		23	64	87	0.57	0.12
(%)	(39.1)	(30.4)	(30.4)		(100.0)				
糖尿病性腎症	30	227	341	145	743	1,583	2,326	0.70	0.12
(%)	(4.0)	(30.6)	(45.9)	(19.5)	(100.0)				
SLE腎炎	2	7	5		14	26	40	0.61	0.12
(%)	(14.3)	(50.0)	(35.7)		(100.0)				
アミロイド腎		2	1	1	4	9	13	0.71	0.17
(%)		(50.0)	(25.0)	(25.0)	(100.0)				
痛風腎	1	4	6	1	12	11	23	0.65	0.11
(%)	(8.3)	(33.3)	(50.0)	(8.3)	(100.0)				
先天性代謝異常による腎不全	1	2	2	1	6	9	15	0.63	0.17
(%)	(16.7)	(33.3)	(33.3)	(16.7)	(100.0)				
腎・尿路結核						2	2		
(%)									
腎・尿路結石	1	1	4	2	8	9	17	0.70	0.16
(%)	(12.5)	(12.5)	(50.0)	(25.0)	(100.0)				
腎・尿路腫瘍	1	1			2	8	10	0.54	0.15
(%)	(50.0)	(50.0)			(100.0)				
閉塞性尿路障害	2	4	3		9	10	19	0.58	0.11
(%)	(22.2)	(44.4)	(33.3)		(100.0)				
骨髄腫			1	1	2	5	7	0.75	0.09
(%)			(50.0)	(50.0)	(100.0)				
腎形成不全	4	8	12	1	25	42	67	0.62	0.17
(%)	(16.0)	(32.0)	(48.0)	(4.0)	(100.0)				
不明	19	77	93	16	205	477	682	0.65	0.13
(%)	(9.3)	(37.6)	(45.4)	(7.8)	(100.0)				
再導入	1	7	10	1	19	22	41	0.66	0.11
(%)	(5.3)	(36.8)	(52.6)	(5.3)	(100.0)				
その他	14	23	38	14	89	142	231	0.66	0.15
(%)	(15.7)	(25.8)	(42.7)	(15.7)	(100.0)				
合計	221	853	1,068	344	2,486	4,618	7,104	0.66	0.13
(%)	(8.9)	(34.3)	(43.0)	(13.8)	(100.0)				
記載なし									
(%)									
総計	221	853	1,068	344	2,486	4,618	7,104	0.66	0.13
(%)	(8.9)	(34.3)	(43.0)	(13.8)	(100.0)				

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査