

表 3782 腹膜炎罹患回数(回/年) 年齢(歳)別 (腹膜透析単独患者)

年齢	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
15歳未満	51	6	2	1							60		29	89
(%)	(85.0)	(10.0)	(3.3)	(1.7)							(100.0)			
15歳～	50	6	1								57		31	88
(%)	(87.7)	(10.5)	(1.8)								(100.0)			
30歳～	264	31	3			1					299	3	201	503
(%)	(88.3)	(10.4)	(1.0)			(0.3)					(100.0)			
45歳～	955	98	19	3	2	1			1		1,079	2	645	1,726
(%)	(88.5)	(9.1)	(1.8)	(0.3)	(0.2)	(0.1)			(0.1)		(100.0)			
60歳～	1,630	215	46	14	4			1			1,910	4	1,163	3,077
(%)	(85.3)	(11.3)	(2.4)	(0.7)	(0.2)			(0.1)			(100.0)			
75歳～	733	123	31	5	3	2	1				898	2	596	1,496
(%)	(81.6)	(13.7)	(3.5)	(0.6)	(0.3)	(0.2)	(0.1)				(100.0)			
90歳～	63	7	3	1							74	1	50	125
(%)	(85.1)	(9.5)	(4.1)	(1.4)							(100.0)			
合計	3,746	486	105	24	9	4	1	1	1		4,377	12	2,715	7,104
(%)	(85.6)	(11.1)	(2.4)	(0.5)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)		(100.0)			
不明														
(%)														
記載なし														
(%)														
総計	3,746	486	105	24	9	4	1	1	1		4,377	12	2,715	7,104
(%)	(85.6)	(11.1)	(2.4)	(0.5)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)		(100.0)			
平均	62.99	64.96	67.32	67.38	66.44	60.25	78.00	65.00	59.00		63.34	62.50	63.77	63.50
標準偏差	15.14	15.04	15.60	16.37	10.08	22.90					15.16	16.26	14.98	15.09

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査