

表 3784 腹膜炎罹患回数(回/年) 透析歴(年)別 (腹膜透析単独患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,867	213	53	10	5	2					2,150	6	1,308	3,464
(%)	(86.8)	(9.9)	(2.5)	(0.5)	(0.2)	(0.1)					(100.0)			
2年～	1,288	197	35	12	2	1	1	1			1,537	4	961	2,502
(%)	(83.8)	(12.8)	(2.3)	(0.8)	(0.1)	(0.1)	(0.1)	(0.1)			(100.0)			
5年～	492	61	14		2	1			1		571	2	350	923
(%)	(86.2)	(10.7)	(2.5)		(0.4)	(0.2)			(0.2)		(100.0)			
10年～	67	7	3	1							78		57	135
(%)	(85.9)	(9.0)	(3.8)	(1.3)							(100.0)			
15年～	19	4									23		18	41
(%)	(82.6)	(17.4)									(100.0)			
20年～	5										5		8	13
(%)	(100.0)										(100.0)			
25年～	3	3									6		1	7
(%)	(50.0)	(50.0)									(100.0)			
30年～	3										3			3
(%)	(100.0)										(100.0)			
35年～				1							1			1
(%)				(100.0)							(100.0)			
40年～	1										1		1	2
(%)	(100.0)										(100.0)			
合計	3,745	485	105	24	9	4	1	1	1		4,375	12	2,704	7,091
(%)	(85.6)	(11.1)	(2.4)	(0.5)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)		(100.0)			
不明	1	1									2		11	13
(%)	(50.0)	(50.0)									(100.0)			
記載なし														
(%)														
総計	3,746	486	105	24	9	4	1	1	1		4,377	12	2,715	7,104
(%)	(85.6)	(11.1)	(2.4)	(0.5)	(0.2)	(0.1)	(0.0)	(0.0)	(0.0)		(100.0)			
平均	2.35	2.60	2.30	3.46	2.56	3.00	2.00	2.00	5.00		2.39	2.00	2.42	2.40
標準偏差	2.98	3.20	2.42	7.05	2.83	3.16					3.03	2.09	3.05	3.04

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査