

表 3822 出口部感染罹患回数(回/年) 透析歴(年)別 (腹膜透析患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,791	328	75	31	12	5	3	2			2,247	8	1,425	3,680
(%)	(79.7)	(14.6)	(3.3)	(1.4)	(0.5)	(0.2)	(0.1)	(0.1)			(100.0)			
2年～	1,481	264	75	29	13	6	3	1	2	4	1,878	6	1,218	3,102
(%)	(78.9)	(14.1)	(4.0)	(1.5)	(0.7)	(0.3)	(0.2)	(0.1)	(0.1)	(0.2)	(100.0)			
5年～	732	118	31	8	5	1			1		896	2	622	1,520
(%)	(81.7)	(13.2)	(3.5)	(0.9)	(0.6)	(0.1)			(0.1)		(100.0)			
10年～	160	20	7	3		1					191		123	314
(%)	(83.8)	(10.5)	(3.7)	(1.6)		(0.5)					(100.0)			
15年～	46	2	1	1							50		37	87
(%)	(92.0)	(4.0)	(2.0)	(2.0)							(100.0)			
20年～	12	3	1								16		10	26
(%)	(75.0)	(18.8)	(6.3)								(100.0)			
25年～	6	2									8		4	12
(%)	(75.0)	(25.0)									(100.0)			
30年～	3										3		3	6
(%)	(100.0)										(100.0)			
35年～	2										2			2
(%)	(100.0)										(100.0)			
40年～	1										1		1	2
(%)	(100.0)										(100.0)			
合計	4,234	737	190	72	30	13	6	3	3	4	5,292	16	3,443	8,751
(%)	(80.0)	(13.9)	(3.6)	(1.4)	(0.6)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(100.0)			
不明	4										4		14	18
(%)	(100.0)										(100.0)			
記載なし														
(%)														
総計	4,238	737	190	72	30	13	6	3	3	4	5,296	16	3,457	8,769
(%)	(80.0)	(13.9)	(3.6)	(1.4)	(0.6)	(0.2)	(0.1)	(0.1)	(0.1)	(0.1)	(100.0)			
平均	3.03	2.80	2.94	2.72	2.20	2.62	1.33	1.00	3.67	2.25	2.98	2.06	3.02	2.99
標準偏差	3.66	3.26	3.12	2.91	1.85	2.93	1.21	1.00	2.08	0.50	3.57	1.91	3.54	3.55

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査