

表 3839 出口部感染罹患回数(回/年) 透析歴(年)別 (腹膜透析単独患者)

透析歴	なし	1回	2回	3回	4回	5回	6回	7回	8回	9回以上	合計	不明	記載なし	総計
2年未満	1,692	308	70	29	11	5	3	2			2,120	7	1,337	3,464
(%)	(79.8)	(14.5)	(3.3)	(1.4)	(0.5)	(0.2)	(0.1)	(0.1)			(100.0)			
2年～	1,202	209	56	26	13	6	3	1	2	2	1,520	4	978	2,502
(%)	(79.1)	(13.8)	(3.7)	(1.7)	(0.9)	(0.4)	(0.2)	(0.1)	(0.1)	(0.1)	(100.0)			
5年～	452	75	22	6	5	1			1		562	2	359	923
(%)	(80.4)	(13.3)	(3.9)	(1.1)	(0.9)	(0.2)			(0.2)		(100.0)			
10年～	65	7	2	2							76		59	135
(%)	(85.5)	(9.2)	(2.6)	(2.6)							(100.0)			
15年～	21	1		1							23		18	41
(%)	(91.3)	(4.3)		(4.3)							(100.0)			
20年～	3	1	1								5		8	13
(%)	(60.0)	(20.0)	(20.0)								(100.0)			
25年～	4	2									6		1	7
(%)	(66.7)	(33.3)									(100.0)			
30年～	3										3			3
(%)	(100.0)										(100.0)			
35年～	1										1			1
(%)	(100.0)										(100.0)			
40年～	1										1		1	2
(%)	(100.0)										(100.0)			
合計	3,444	603	151	64	29	12	6	3	3	2	4,317	13	2,761	7,091
(%)	(79.8)	(14.0)	(3.5)	(1.5)	(0.7)	(0.3)	(0.1)	(0.1)	(0.1)	(0.0)	(100.0)			
不明	2										2		11	13
(%)	(100.0)										(100.0)			
記載なし														
(%)														
総計	3,446	603	151	64	29	12	6	3	3	2	4,319	13	2,772	7,104
(%)	(79.8)	(14.0)	(3.5)	(1.5)	(0.7)	(0.3)	(0.1)	(0.1)	(0.1)	(0.0)	(100.0)			
平均	2.40	2.32	2.46	2.52	2.24	2.00	1.33	1.00	3.67	2.50	2.39	2.08	2.42	2.40
標準偏差	3.10	2.88	2.70	2.82	1.86	2.00	1.21	1.00	2.08	0.71	3.04	1.98	3.04	3.04

数値下のかっこ内は行方向の合計に対する%です。

(C)Japanese Society for Dialysis Therapy

2015年調査